

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C12N 15/82, 15/79, 15/54, A01H 5/00		A2	(11) International Publication Number: WO 97/04113
			(43) International Publication Date: 6 February 1997 (06.02.97)
(21) International Application Number: PCT/EP96/03053 (22) International Filing Date: 12 July 1996 (12.07.96) (30) Priority Data: 9514437.4 14 July 1995 (14.07.95) GB (71) Applicant (for all designated States except US): DANISCO A/S [DK/DK]; Langebrogade 1, P.O. Box 17, DK-1001 Copenhagen K (DK). (72) Inventor; and (75) Inventor/Applicant (for US only): POULSEN, Peter [DK/DK]; Holsteinsgade 52, 3, DK-2100 Copenhagen K (DK). (74) Agents: HARDING, Charles, Thomas et al.; D. Young & Co., 21 New Fetter Lane, London EC4A 1DA (GB).			(81) Designated States: AL, AM, AT, AU, AZ, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, EE, ES, FI, GB, GE, HU, IL, IS, JP, KE, KG, KP, KR, KZ, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TR, TT, UA, UG, US, UZ, VN, ARIPO patent (KE, LS, MW, SD, SZ, UG), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published <i>Without international search report and to be republished upon receipt of that report. With an indication in relation to a deposited microorganism furnished under Rule 13^{bis} separately from the description. Date of receipt by the International Bureau: 20 August 1996 (20.08.96)</i>
(54) Title: INHIBITION OF GENE EXPRESSION			
(57) Abstract A method of inhibiting gene expression is described. The method, which affects enzymatic activity in a plant, comprises expressing in a plant (or a cell, a tissue or an organ thereof) a nucleotide sequence wherein the nucleotide sequence codes, partially or completely, for an intron in a sense orientation; and wherein the nucleotide sequence does not contain a sequence that is sense to an exon sequence normally associated with the intron.			

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AM	Armenia	GB	United Kingdom	MW	Malawi
AT	Austria	GE	Georgia	MX	Mexico
AU	Australia	GN	Guinea	NE	Niger
BB	Barbados	GR	Greece	NL	Netherlands
BE	Belgium	HU	Hungary	NO	Norway
BF	Burkina Faso	IE	Ireland	NZ	New Zealand
BG	Bulgaria	IT	Italy	PL	Poland
BJ	Benin	JP	Japan	PT	Portugal
BR	Brazil	KE	Kenya	RO	Romania
BY	Belarus	KG	Kyrgyzstan	RU	Russian Federation
CA	Canada	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	KZ	Kazakhstan	SG	Singapore
CH	Switzerland	LI	Liechtenstein	SI	Slovenia
CI	Côte d'Ivoire	LK	Sri Lanka	SK	Slovakia
CM	Cameroon	LR	Liberia	SN	Senegal
CN	China	LT	Lithuania	SZ	Swaziland
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	LV	Latvia	TG	Togo
DE	Germany	MC	Monaco	TJ	Tajikistan
DK	Denmark	MD	Republic of Moldova	TT	Trinidad and Tobago
EE	Estonia	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	UG	Uganda
FI	Finland	MN	Mongolia	US	United States of America
FR	France	MR	Mauritania	UZ	Uzbekistan
GA	Gabon			VN	Viet Nam

INHIBITION OF GENE EXPRESSION

5 The present invention relates to a method of inhibiting gene expression, particularly inhibiting gene expression in a plant. The present invention also relates to a nucleotide sequence useful in the method. In addition, the present invention relates to a promoter that is useful for expressing the nucleotide sequence.

10 Starch is one of the main storage carbohydrates in plants, especially higher plants. The structure of starch consists of amylose and amylopectin. Amylose consists essentially of straight chains of α -1-4-linked glycosyl residues. Amylopectin comprises chains of α -1-4-linked glycosyl residues with some α -1-6 branches. The branched nature of amylopectin is accomplished by the action of *inter alia* an enzyme commonly known as the starch branching enzyme ("SBE"). SBE catalyses the formation of branch points in the amylopectin molecule by adding α -1,4 glucans through α -1,6-glucosidic branching linkages. The biosynthesis of amylose and amylopectin is schematically shown in Figure 1, whereas the α -1-4-links and the α -1-6 links are shown in Figure 2.

20 It is known that starch is an important raw material. Starch is widely used in the food, paper, and chemical industries. However, a large fraction of the starches used in these industrial applications are post-harvest modified by chemical, physical or enzymatic methods in order to obtain starches with certain required functional properties.

25 Within the past few years it has become desirable to make genetically modified plants which could be capable of producing modified starches which could be the same as the post-harvest modified starches. It is also known that it may be possible to prepare such genetically modified plants by expression of antisense nucleotide coding sequences. In this regard, June Bourque provides a detailed summary of antisense strategies for the genetic manipulations in plants (Bourque 1995 Plant Science 105 pp 30 125-149).

Whilst it is known that enzymatic activity can be affected by expression of particular nucleotide sequences (for example see the teachings of Finnegan and McElroy [1994] Biotechnology 12 883-888; and Matzke and Matzke [1995] TIG 11 1-3) there is still a need for a method that can more reliably and/or more efficiently and/or more specifically affect enzymatic activity.

According to a first aspect of the present invention there is provided a method of affecting enzymatic activity in a plant (or a cell, a tissue or an organ thereof) comprising expressing in the plant (or a cell, a tissue or an organ thereof) a nucleotide sequence wherein the nucleotide sequence partially or completely codes (is) an intron in a sense orientation; and wherein the nucleotide sequence does not contain a sequence that is a sense exon sequence normally associated with the intron.

According to a second aspect of the present invention there is provided a method of affecting enzymatic activity in a starch producing organism (or a cell, a tissue or an organ thereof) comprising expressing in the starch producing organism (or a cell, a tissue or an organ thereof) a nucleotide sequence wherein the nucleotide sequence codes, partially or completely, for an intron in a sense orientation; wherein the nucleotide sequence does not contain a sequence that is sense to an exon sequence normally associated with the intron; and wherein starch branching enzyme activity is affected and/or the levels of amylopectin are affected and/or the composition of starch is changed.

According to a third aspect of the present invention there is provided a sequence comprising the nucleotide sequence shown as any one of SEQ.I.D. No. 1 to SEQ.I.D. No. 13 or a variant, derivative or homologue thereof.

According to a fourth aspect of the present invention there is provided a promoter comprising the sequence shown as SEQ.I.D. No. 14 or a variant, derivative or homologue thereof.

According to a fifth aspect of the present invention there is provided a construct capable of comprising or expressing the present invention.

5 According to a sixth aspect of the present invention there is provided a vector comprising or expressing the present invention.

According to a seventh aspect of the present invention there is provided a cell, tissue or organ comprising or expressing the present invention.

10 According to an eighth aspect of the present invention there is provided a transgenic starch producing organism comprising or expressing the present invention. According to a ninth aspect of the present invention there is provided a starch obtained from the present invention.

15 According to a tenth aspect of the present invention there is provided pBEA11 (NCIMB 40754). According to an eleventh aspect of the present invention there is provided a sense nucleotide sequence that is obtainable from λ -SBE 3.2 (NCIMB 40751) or λ -SBE 3.4 (NCIMB 40752) or a variant, derivative or homologue thereof.

20 A key advantage of the present invention is that it provides a method for preparing modified starches that is not dependent on the need for post-harvest modification of starches. Thus the method of the present invention obviates the need for the use of hazardous chemicals that are normally used in the post-harvest modification of starches.

25 In addition, the present invention provides *inter alia* genetically modified plants which are capable of producing modified and/or novel and/or improved starches whose properties would satisfy various industrial requirements.

30 Thus, the present invention provides a method of preparing tailor-made starches in plants which could replace the post-harvest modified starches.

Also, the present invention provides a method that enables modified starches to be prepared by a method that can have a more beneficial effect on the environment than the known post-harvest modification methods which are dependent on the use of hazardous chemicals and large quantities of energy.

5

An other key advantage of the present invention is that it provides a method that may more reliably and/or more efficiently and/or more specifically affect enzymatic activity when compared to the known methods of affecting enzymatic activity. With regard to this advantage of the present invention it is to be noted that there is some degree of homology between coding regions of SBEs. However, there is little or no homology with the intron sequences of SBEs. Thus, sense intron expression provides a mechanism to affect selectively the expression of a particular SBE. This advantageous aspect could be used, for example, to reduce or eliminate a particular SBE enzyme and replace that enzyme with another enzyme which can be another branching enzyme or even a recombinant version of the affected enzyme or even a hybrid enzyme which could for example comprise part of a SBE enzyme from one source and at least a part of another SBE enzyme from another source. This particular feature of the present invention is covered by the combination aspect of the present invention which is discussed in more detail later.

10

15

20

Thus the present invention provides a mechanism for selectively affecting SBE activity. This is in contrast to the prior art methods which are dependent on the use of for example antisense exon expression whereby it would not be possible to introduce new SBE activity without affecting that activity as well.

25

Preferably with the first aspect of the present invention starch branching enzyme activity is affected and/or wherein the levels of amylopectin are affected and/or the composition of starch is changed.

30

Preferably with the first or second aspect of the present invention the nucleotide sequence does not contain a sequence that is sense to an exon sequence.

Preferably with the fourth aspect of the present invention the promoter is in combination with a gene of interest ("GOI").

Preferably the enzymatic activity is reduced or eliminated.

5

Preferably the nucleotide sequence codes for at least substantially all of at least one intron in a sense orientation.

10

Preferably the nucleotide sequence codes, partially or completely, for two or more introns and wherein each intron is in a sense orientation.

Preferably the nucleotide sequence comprises at least 350 nucleotides (e.g. 350 bp), more preferably at least 500 nucleotides (e.g. 500 bp).

15

Preferably the nucleotide sequence comprises the sequence shown as any one of SEQ. I.D. No. 1 to SEQ.I.D. No. 13 or a variant, derivative or homologue thereof, including combinations thereof.

20

Preferably the nucleotide sequence is expressed by a promoter having a sequence shown as SEQ. I.D. No. 14 or a variant, derivative or homologue thereof.

Preferably the transgenic starch producing organism is a plant.

25

A preferred aspect of the present invention therefore relates to a method of affecting enzymatic activity in a plant (or a cell, a tissue or an organ thereof) comprising expressing in the plant (or a cell, a tissue or an organ thereof) a nucleotide sequence wherein the nucleotide sequence codes, partially or completely, for an intron in a sense orientation; wherein the nucleotide sequence does not contain a sequence that is sense to an exon sequence normally associated with the intron; and wherein starch branching enzyme activity is affected and/or the levels of amylopectin are affected and/or the composition of starch is changed.

30

A more preferred aspect of the present invention therefore relates to a method of affecting enzymatic activity in a plant (or a cell, a tissue or an organ thereof) comprising expressing in the plant (or a cell, a tissue or an organ thereof) a nucleotide sequence wherein the nucleotide sequence codes, partially or completely, for an intron in a sense orientation; wherein the nucleotide sequence does not contain a sequence that is sense to an exon sequence normally associated with the intron; wherein starch branching enzyme activity is affected and/or the levels of amylopectin are affected and/or the composition of starch is changed; and wherein the nucleotide sequence comprises the sequence shown as any one of SEQ.I.D. No. 1 to SEQ.I.D. No. 13 or a variant, derivative or homologue thereof, including combinations thereof.

The term "nucleotide" in relation to the present invention includes DNA and RNA. Preferably it means DNA, more preferably DNA prepared by use of recombinant DNA techniques.

The term "intron" is used in its normal sense as meaning a segment of nucleotides, usually DNA, that does not encode part or all of an expressed protein or enzyme.

The term "exon" is used in its normal sense as meaning a segment of nucleotides, usually DNA, encoding part or all of an expressed protein or enzyme.

Thus, the term "intron" refers to gene regions that are transcribed into RNA molecules, but which are spliced out of the RNA before the RNA is translated into a protein. In contrast, the term "exon" refers to gene regions that are transcribed into RNA and subsequently translated into proteins.

The terms "variant" or "homologue" or "fragment" in relation to the nucleotide sequence of the present invention include any substitution of, variation of, modification of, replacement of, deletion of or addition of one (or more) nucleic acid from or to the respective nucleotide sequence providing the resultant nucleotide sequence can affect enzyme activity in a plant, or cell or tissue thereof, preferably wherein the resultant nucleotide sequence has at least the same effect as any one of

the sense sequences shown as SEQ.I.D. No.s 1-13. In particular, the term "homologue" covers homology with respect to similarity of structure and/or similarity of function providing the resultant nucleotide sequence has the ability to affect enzymatic activity in accordance with the present invention. With respect to sequence homology (i.e. similarity), preferably there is more than 80% homology, more preferably at least 85% homology, more preferably at least 90% homology, even more preferably at least 95% homology, more preferably at least 98% homology. The above terms are also synonymous with allelic variations of the sequences.

10 Likewise, the terms "variant" or "homologue" or "fragment" in relation to the promoter of the present invention include any substitution of, variation of, modification of, replacement of, deletion of or addition of one (or more) nucleic acid from or to the respective promoter sequence providing the resultant promoter sequence allows expression of a GOI, preferably wherein the resultant promoter sequence has at least the same effect as SEQ.I.D. No. 14. In particular, the term "homologue" covers homology with respect to similarity of structure and/or similarity of function providing the resultant promoter sequence has the ability to allow for expression of a GOI, such as a nucleotide sequence according to the present invention. With respect to sequence homology (i.e. similarity), preferably there is more than 80% homology, more preferably at least 85% homology, more preferably at least 90% homology, even more preferably at least 95% homology, more preferably at least 98% homology. The above terms are also synonymous with allelic variations of the sequences.

25 The intron sequence of the present invention can be any one or all of the intron sequences of the present invention, including partial sequences thereof, provided that if partial sense sequences are used (i.e. sequences that are not or do not comprise any one or more of the full sequences shown as SEQ.I.D. No.1-13) the partial sequences affect enzymatic activity. Suitable examples of partial sequences include sequences that are shorter than any one of the full sense sequences shown as SEQ.I.D.No.s 1 to 13 but which comprise nucleotides that are adjacent the respective exon or exons.

With regard to the second aspect of the present invention (i.e. specifically affecting SBE activity), the nucleotide sequences of the present invention may comprise one or more sense or antisense exon sequences of the SBE gene (but not sense exon sequences naturally associated with the intron sequence), including complete or partial sequences thereof, providing the nucleotide sequences can affect SBE activity, preferably wherein the nucleotide sequences reduce or eliminate SBE activity. Preferably, the nucleotide sequence of the second aspect of the present invention does not comprise sense exon sequences.

The term "vector" includes an expression vector and a transformation vector. The term "expression vector" means a construct capable of *in vivo* or *in vitro* expression. The term "transformation vector" means a construct capable of being transferred from one species to another - such as from an *E.Coli* plasmid to a fungus or a plant cell, or from an *Agrobacterium* to a plant cell.

The term "construct" - which is synonymous with terms such as "conjugate", "cassette" and "hybrid" - in relation to the sense nucleotide sequence aspect of the present invention includes the nucleotide sequence according to the present invention directly or indirectly attached to a promoter. An example of an indirect attachment is the provision of a suitable spacer group such as an intron sequence, such as the *Shl*-intron or the ADH intron, intermediate the promoter and the nucleotide sequence of the present invention. The same is true for the term "fused" in relation to the present invention which includes direct or indirect attachment. The terms do not cover the natural combination of the wild type SBE gene when associated with the wild type SBE gene promoter in their natural environment.

The construct may even contain or express a marker which allows for the selection of the genetic construct in, for example, a plant cell into which it has been transferred. Various markers exist which may be used in, for example, plants - such as mannose. Other examples of markers include those that provide for antibiotic resistance - e.g. resistance to G418, hygromycin, bleomycin, kanamycin and gentamycin.

The construct of the present invention preferably comprises a promoter. The term "promoter" is used in the normal sense of the art, e.g. an RNA polymerase binding site in the Jacob-Monod theory of gene expression. Examples of suitable promoters are those that can direct efficient expression of the nucleotide sequence of the present invention and/or in a specific type of cell. Some examples of tissue specific promoters are disclosed in WO 92/11375.

The promoter could additionally include conserved regions such as a Pribnow Box or a TATA box. The promoters may even contain other sequences to affect (such as to maintain, enhance, decrease) the levels of expression of the nucleotide sequence of the present invention. Suitable examples of such sequences include the *Shl*-intron or an ADH intron. Other sequences include inducible elements - such as temperature, chemical, light or stress inducible elements. Also, suitable elements to enhance transcription or translation may be present. An example of the latter element is the TMV 5' leader sequence (see Sleat Gene 217 [1987] 217-225; and Dawson Plant Mol. Biol. 23 [1993] 97).

As mentioned, the construct and/or the vector of the present invention may include a transcriptional initiation region which may provide for regulated or constitutive expression. Any suitable promoter may be used for the transcriptional initiation region, such as a tissue specific promoter. In one aspect, preferably the promoter is the patatin promoter or the E35S promoter. In another aspect, preferably the promoter is the SBE promoter.

If, for example, the organism is a plant then the promoter can be one that affects expression of the nucleotide sequence in any one or more of seed, tuber, stem, sprout, root and leaf tissues, preferably tuber. By way of example, the promoter for the nucleotide sequence of the present invention can be the α -Amy 1 promoter (otherwise known as the Amy 1 promoter, the Amy 637 promoter or the α -Amy 637 promoter) as described in our co-pending UK patent application No. 9421292.5 filed 21 October 1994. Alternatively, the promoter for the nucleotide sequence of the present invention can be the α -Amy 3 promoter (otherwise known as the Amy 3

promoter, the Amy 351 promoter or the α -Amy 351 promoter) as described in our co-pending UK patent application No. 9421286.7 filed 21 October 1994.

5 The present invention also encompasses the use of a promoter to express a nucleotide sequence according to the present invention, wherein a part of the promoter is inactivated but wherein the promoter can still function as a promoter. Partial inactivation of a promoter in some instances is advantageous. In particular, with the Amy 351 promoter mentioned earlier it is possible to inactivate a part of it so that the partially inactivated promoter expresses the nucleotide sequence of the present
10 invention in a more specific manner such as in just one specific tissue type or organ. The term "inactivated" means partial inactivation in the sense that the expression pattern of the promoter is modified but wherein the partially inactivated promoter still functions as a promoter. However, as mentioned above, the modified promoter is capable of expressing a gene coding for the enzyme of the present invention in at least one (but not all) specific tissue of the original promoter. Examples of partial
15 inactivation include altering the folding pattern of the promoter sequence, or binding species to parts of the nucleotide sequence, so that a part of the nucleotide sequence is not recognised by, for example, RNA polymerase. Another, and preferable, way of partially inactivating the promoter is to truncate it to form fragments thereof. Another way would be to mutate at least a part of the sequence so that the RNA
20 polymerase can not bind to that part or another part. Another modification is to mutate the binding sites for regulatory proteins for example the CreA protein known from filamentous fungi to exert carbon catabolite repression, and thus abolish the catabolite repression of the native promoter.

25 The construct and/or the vector of the present invention may include a transcriptional termination region.

30 The nucleotide according to the present invention can be expressed in combination (but not necessarily at the same time) with an additional construct. Thus the present invention also provides a combination of constructs comprising a first construct comprising the nucleotide sequence according to the present invention operatively

linked to a first promoter; and a second construct comprising a GOI operatively linked to a second promoter (which need not be the same as the first promoter). With this aspect of the present invention the combination of constructs may be present in the same vector, plasmid, cells, tissue, organ or organism. This aspect of the present invention also covers methods of expressing the same, preferably in specific cells or tissues, such as expression in just a specific cell or tissue, of an organism, typically a plant. With this aspect of the present invention the second construct does not cover the natural combination of the gene coding for an enzyme ordinarily associated with the wild type gene promoter when they are both in their natural environment.

An example of a suitable combination would be a first construct comprising the nucleotide sequence of the present invention and a promoter, such as the promoter of the present invention, and a second construct comprising a promoter, such as the promoter of the present invention, and a GOI wherein the GOI codes for another starch branching enzyme either in sense or antisense orientation.

The above comments relating to the term "construct" for the sense nucleotide aspect of the present invention are equally applicable to the term "construct" for the promoter aspect of the present invention. In this regard, the term includes the promoter according to the present invention directly or indirectly attached to a GOI.

The term "GOI" with reference to the promoter aspect of the present invention or the combination aspect of the present invention means any gene of interest, which need not necessarily code for a protein or an enzyme - as is explained later. A GOI can be any nucleotide sequence that is either foreign or natural to the organism in question, for example a plant.

Typical examples of a GOI include genes encoding for other proteins or enzymes that modify metabolic and catabolic processes. The GOI may code for an agent for introducing or increasing pathogen resistance.

The GOI may even be an antisense construct for modifying the expression of natural transcripts present in the relevant tissues. An example of such a GOI is the nucleotide sequence according to the present invention.

5 The GOI may even code for a protein that is non-natural to the host organism - e.g. a plant. The GOI may code for a compound that is of benefit to animals or humans. For example, the GOI could code for a pharmaceutically active protein or enzyme such as any one of the therapeutic compounds insulin, interferon, human serum albumin, human growth factor and blood clotting factors. The GOI may even code
10 for a protein giving additional nutritional value to a food or feed or crop. Typical examples include plant proteins that can inhibit the formation of anti-nutritive factors and plant proteins that have a more desirable amino acid composition (e.g. a higher lysine content than a non-transgenic plant). The GOI may even code for an enzyme that can be used in food processing such as xylanases and α -galactosidase. The GOI
15 can be a gene encoding for any one of a pest toxin, an antisense transcript such as that for α -amylase, a protease or a glucanase. Alternatively, the GOI can be a nucleotide sequence according to the present invention.

The GOI can be the nucleotide sequence coding for the arabinofuranosidase enzyme
20 which is the subject of our co-pending UK patent application 9505479.7. The GOI can be the nucleotide sequence coding for the glucanase enzyme which is the subject of our co-pending UK patent application 9505475.5. The GOI can be the nucleotide sequence coding for the α -amylase enzyme which is the subject of our co-pending UK patent application 9413439.2. The GOI can be the nucleotide sequence coding for
25 the α -amylase enzyme which is the subject of our co-pending UK patent application 9421290.9. The GOI can be any of the nucleotide sequences coding for the α -glucan lyase enzyme which are described in our co-pending PCT patent application PCT/EP94/03397.

30 In one aspect the GOI can even be a nucleotide sequence according to the present invention but when operatively linked to a different promoter.

The GOI could include a sequence that codes for one or more of a xylanase, an arabinase, an acetyl esterase, a rhamnogalacturonase, a glucanase, a pectinase, a branching enzyme or another carbohydrate modifying enzyme or proteinase. Alternatively, the GOI may be a sequence that is antisense to any of those sequences.

5

As mentioned above, the present invention provides a mechanism for selectively affecting a particular enzymatic activity.

10 In an important application of the present invention it is now possible to reduce or eliminate expression of a genomic nucleotide sequence coding for a genomic protein or enzyme by expressing a sense intron construct for that particular genomic protein or enzyme and (e.g. at the same time) expressing a recombinant version of that enzyme or protein - in other words the GOI is a recombinant nucleotide sequence coding for the genomic enzyme or protein. This application allows expression of
15 desired recombinant enzymes and proteins in the absence of (or reduced levels of) respective genomic enzymes and proteins. Thus the desired recombinant enzymes and proteins can be easily separated and purified from the host organism. This particular aspect of the present invention is very advantageous over the prior art methods which, for example, rely on the use of anti-sense exon expression which methods also affect
20 expression of the recombinant enzyme.

Thus, a further aspect of the present invention relates to a method of expressing a recombinant protein or enzyme in a host organism comprising expressing a nucleotide sequence coding for the recombinant protein or enzyme; and expressing a further
25 nucleotide sequence wherein the further nucleotide sequence codes, partially or completely, for an intron in a sense orientation; wherein the intron is an intron normally associated with the genomic gene encoding a protein or an enzyme corresponding to the recombinant protein or enzyme; and wherein the further nucleotide sequence does not contain a sequence that is sense to an exon sequence
30 normally associated with the intron. Additional aspects cover the combination of those nucleotide sequences including their incorporation in constructs, vectors, cells, tissues and transgenic organisms.

Therefore the present invention also relates to a combination of nucleotide sequences comprising a first nucleotide sequence coding for a recombinant enzyme; and a second nucleotide sequence which corresponds to an intron in a sense orientation; wherein the intron is an intron that is associated with a genomic gene encoding the enzyme corresponding to the recombinant enzyme; and wherein the second nucleotide sequence does not contain a sequence that is sense to an exon sequence normally associated with the intron.

The GOI may even code for one or more introns but in an antisense orientation, such as any one or more of the antisense intron sequences presented in the attached sequence listings. For example, the present invention also covers the expression of for example a sense intron (e.g. SEQ.I.D.No. 1) in combination with for example an antisense sense intron which preferably is not complementary to the sense intron sequence (e.g. SEQ.I.D.No. 16).

The terms "cell", "tissue" and "organ" include cell, tissue and organ *per se* and when within an organism.

The term "organism" in relation to the present invention includes any organism that could comprise the nucleotide sequence according to the present invention and/or wherein the nucleotide sequence according to the present invention can be expressed when present in the organism. Preferably the organism is a starch producing organism such as any one of a plant, algae, fungi, yeast and bacteria, as well as cell lines thereof. Preferably the organism is a plant.

The term "starch producing organism" includes any organism that can biosynthesise starch. Preferably, the starch producing organism is a plant.

The term "plant" as used herein includes any suitable angiosperm, gymnosperm, monocotyledon and dicotyledon. Typical examples of suitable plants include vegetables such as potatoes; cereals such as wheat, maize, and barley; fruit; trees; flowers; and other plant crops. Preferably, the term means "potato".

The term "transgenic organism" in relation to the present invention includes any organism that comprises the nucleotide sequence according to the present invention and/or products obtained therefrom, and/or wherein the nucleotide sequence according to the present invention can be expressed within the organism. Preferably the nucleotide sequence of the present invention is incorporated in the genome of the organism. Preferably the transgenic organism is a plant, more preferably a potato.

To prepare the host organism one can use prokaryotic or eukaryotic organisms. Examples of suitable prokaryotic hosts include *E. coli* and *Bacillus subtilis*. Teachings on the transformation of prokaryotic hosts is well documented in the art, for example see Sambrook *et al* (Sambrook *et al.* in Molecular Cloning: A Laboratory Manual, 2nd edition, 1989, Cold Spring Harbor Laboratory Press).

Even though the enzyme according to the present invention and the nucleotide sequence coding for same are not disclosed in EP-B-0470145 and CA-A-2006454, those two documents do provide some useful background commentary on the types of techniques that may be employed to prepare transgenic plants according to the present invention. Some of these background teachings are now included in the following commentary.

The basic principle in the construction of genetically modified plants is to insert genetic information in the plant genome so as to obtain a stable maintenance of the inserted genetic material.

Several techniques exist for inserting the genetic information, the two main principles being direct introduction of the genetic information and introduction of the genetic information by use of a vector system. A review of the general techniques may be found in articles by Potrykus (Annu Rev Plant Physiol Plant Mol Biol [1991] 42:205-225) and Christou (Agro-Food-Industry Hi-Tech March/April 1994 17-27).

Thus, in one aspect, the present invention relates to a vector system which carries a nucleotide sequence or construct according to the present invention and which is

capable of introducing the nucleotide sequence or construct into the genome of an organism, such as a plant.

5 The vector system may comprise one vector, but it can comprise two vectors. In the case of two vectors, the vector system is normally referred to as a binary vector system. Binary vector systems are described in further detail in Gynheung An *et al.* (1980), Binary Vectors, *Plant Molecular Biology Manual A3*, 1-19.

10 One extensively employed system for transformation of plant cells with a given promoter or nucleotide sequence or construct is based on the use of a Ti plasmid from *Agrobacterium tumefaciens* or a Ri plasmid from *Agrobacterium rhizogenes* An *et al.* (1986), *Plant Physiol.* 81, 301-305 and Butcher D.N. *et al.* (1980), *Tissue Culture Methods for Plant Pathologists*, eds.: D.S. Ingrams and J.P. Helgeson, 203-208. Several different Ti and Ri plasmids have been constructed which are suitable for the
15 construction of the plant or plant cell constructs described above. A non-limiting example of such a Ti plasmid is pGV3850.

The nucleotide sequence or construct of the present invention should preferably be inserted into the Ti-plasmid between the terminal sequences of the T-DNA or adjacent
20 a T-DNA sequence so as to avoid disruption of the sequences immediately surrounding the T-DNA borders, as at least one of these regions appears to be essential for insertion of modified T-DNA into the plant genome.

As will be understood from the above explanation, if the organism is a plant the
25 vector system of the present invention is preferably one which contains the sequences necessary to infect the plant (e.g. the *vir* region) and at least one border part of a T-DNA sequence, the border part being located on the same vector as the genetic construct.

30 Furthermore, the vector system is preferably an *Agrobacterium tumefaciens* Ti-plasmid or an *Agrobacterium rhizogenes* Ri-plasmid or a derivative thereof. As these plasmids are well-known and widely employed in the construction of transgenic

plants, many vector systems exist which are based on these plasmids or derivatives thereof.

5 In the construction of a transgenic plant the nucleotide sequence or construct of the present invention may be first constructed in a microorganism in which the vector can replicate and which is easy to manipulate before insertion into the plant. An example of a useful microorganism is *E. coli*, but other microorganisms having the above properties may be used. When a vector of a vector system as defined above has been constructed in *E. coli*, it is transferred, if necessary, into a suitable *Agrobacterium* 10 strain, e.g. *Agrobacterium tumefaciens*. The Ti-plasmid harbouring the nucleotide sequence or construct of the present invention is thus preferably transferred into a suitable *Agrobacterium* strain, e.g. *A. tumefaciens*, so as to obtain an *Agrobacterium* cell harbouring the promoter or nucleotide sequence or construct of the present invention, which DNA is subsequently transferred into the plant cell to be modified.

15 If, for example, for the transformation the Ti- or Ri-plasmid of the plant cells is used, at least the right boundary and often however the right and the left boundary of the Ti- and Ri-plasmid T-DNA, as flanking areas of the introduced genes, can be connected. The use of T-DNA for the transformation of plant cells has been 20 intensively studied and is described in EP-A-120516; Hoekema, in: The Binary Plant Vector System Offset-drukkerij Kanter B.B., Alblasterdam, 1985, Chapter V; Fraley, *et al.*, Crit. Rev. Plant Sci., 4:1-46; and An *et al.*, EMBO J. (1985) 4:277-284.

25 Direct infection of plant tissues by *Agrobacterium* is a simple technique which has been widely employed and which is described in Butcher D.N. *et al.* (1980), *Tissue Culture Methods for Plant Pathologists*, eds.: D.S. Ingrams and J.P. Helgeson, 203-208. For further teachings on this topic see Potrykus (Annu Rev Plant Physiol Plant-Mol Biol [1991] 42:205-225) and Christou (Agro-Food-Industry Hi-Tech March/April 30 1994 17-27). With this technique, infection of a plant may be performed in or on a certain part or tissue of the plant, i.e. on a part of a leaf, a root, a stem or another part of the plant.

Typically, with direct infection of plant tissues by *Agrobacterium* carrying the GOI (such as the nucleotide sequence according to the present invention) and, optionally, a promoter, a plant to be infected is wounded, e.g. by cutting the plant with a razor blade or puncturing the plant with a needle or rubbing the plant with an abrasive.

5 The wound is then inoculated with the *Agrobacterium*. The inoculated plant or plant part is then grown on a suitable culture medium and allowed to develop into mature plants.

10 When plant cells are constructed, these cells may be grown and maintained in accordance with well-known tissue culturing methods such as by culturing the cells in a suitable culture medium supplied with the necessary growth factors such as amino acids, plant hormones, vitamins, etc.

15 Regeneration of the transformed cells into genetically modified plants may be accomplished using known methods for the regeneration of plants from cell or tissue cultures, for example by selecting transformed shoots using an antibiotic and by subculturing the shoots on a medium containing the appropriate nutrients, plant hormones, etc.

20 Further teachings on plant transformation may be found in EP-A-0449375.

As reported in CA-A-2006454, a large amount of cloning vectors are available which contain a replication system in *E. coli* and a marker which allows a selection of the transformed cells. The vectors contain for example pBR 322, pUC series, M13 mp series, pACYC 184 etc. In this way, the nucleotide or construct of the present invention can be introduced into a suitable restriction position in the vector. The contained plasmid is then used for the transformation in *E. coli*. The *E. coli* cells are cultivated in a suitable nutrient medium and then harvested and lysed. The plasmid is then recovered. As a method of analysis there is generally used sequence analysis, 25 restriction analysis, electrophoresis and further biochemical-molecular biological methods. After each manipulation, the used DNA sequence can be restricted and connected with the next DNA sequence. Each sequence can be cloned in the same 30

or different plasmid.

After the introduction of the nucleotide sequence or construct according to the present invention in the plants the presence and/or insertion of further DNA sequences may
5 be necessary - such as to create combination systems as outlined above (e.g. an organism comprising a combination of constructs).

The above commentary for the transformation of prokaryotic organisms and plants with the nucleotide sequence of the present invention is equally applicable for the
10 transformation of those organisms with the promoter of the present invention.

In summation, the present invention relates to affecting enzyme activity by expressing sense intron sequences.

15 Also, the present invention relates to a promoter useful for the expression of those sense intron sequences.

The following samples have been deposited in accordance with the Budapest Treaty at the recognised depositary The National Collections of Industrial and Marine
20 Bacteria Limited (NCIMB) at 23 St Machar Drive, Aberdeen, Scotland, AB2 1RY, United Kingdom, on 13 July 1995:

NCIMB 40754 (which refers to pBEA 11 as described herein);

25 NCIMB 40751 (which refers to λ -SBE 3.2 as described herein), and

NCIMB 40752 (which refers to λ -SBE 3.4 as described herein).

A highly preferred embodiment of the present invention therefore relates to a method
30 of affecting enzymatic activity in a plant (or a cell, a tissue or an organ thereof) comprising expressing in the plant (or a cell, a tissue or an organ thereof) a nucleotide sequence wherein the nucleotide sequence codes, partially or completely,

for an intron in a sense orientation; wherein the nucleotide sequence does not contain a sequence that is sense to an exon sequence normally associated with the intron; wherein starch branching enzyme activity is affected and/or the levels of amylopectin are affected and/or the composition of starch is changed; and wherein the intron nucleotide sequence is obtainable from NCIMB 40751, NCIMB 40752, or NCIMB 40754 or a variant, derivative or homologue thereof.

A more highly preferred aspect of the present invention therefore relates to a method of affecting enzymatic activity in a plant (or a cell, a tissue or an organ thereof) comprising expressing in the plant (or a cell, a tissue or an organ thereof) a nucleotide sequence wherein the nucleotide sequence codes, partially or completely, for an intron in a sense orientation; wherein the nucleotide sequence does not contain a sequence that is sense to an exon sequence normally associated with the intron; wherein starch branching enzyme activity is affected and/or the levels of amylopectin are affected and/or the composition of starch is changed; wherein the nucleotide sequence comprises the sequence shown as any one of SEQ.I.D. No. 1 to SEQ.I.D. No. 13 or a variant, derivative or homologue thereof, including combinations thereof; and wherein the intron nucleotide sequence is obtainable from NCIMB 40751, NCIMB 40752, or NCIMB 40754, or a variant, derivative or homologue thereof.

The present invention will now be described only by way of example, in which reference is made to the following attached Figures:

Figure 1, which is a schematic representation of the biosynthesis of amylose and amylopectin;

Figure 2, which is a diagrammatic representation of the α -1-4-links and the α -1-6 links of amylopectin;

Figure 3, which is a diagrammatic representation of the exon-intron structure of a genomic SBE clone;

Figure 4, which is a plasmid map of pPATA1, which is 3936 bp in size;

Figure 5, which is a plasmid map of pABE7, which is 5106 bp in size;

5 Figure 6, which is a plasmid map of pVictorIV Man, which is 7080 bp in size;

Figure 7, which is a plasmid map of pBEA11, which is 9.54 kb in size;

10 Figure 8, which shows the full genomic nucleotide sequence for SBE including the promoter, exons and introns;

Figure 9, which is a plasmid map of pVictor5a, which is 9.12 kb in size; and

15 Figure 10, which is a plasmid map of pBEP2, which is 10.32 kb in size.

Figures 1 and 2 were referred to above in the introductory description concerning starch in general. As mentioned, Figure 3 is a diagrammatic representation of the exon-intron structure of a genomic SBE clone, the sequence of which is shown in Figure 8. This clone, which has about 11.5 k base pairs, comprises 14 exons and 13
20 introns. The introns are numbered in increasing order from the 5' end to the 3' end and correspond to SEQ.I.D.No.s 1-13, respectively. Their respective antisense intron sequences are shown as SEQ.I.D.No.s 15-27.

In more detail, Figures 3 and 8 present information on the 11468 base pairs of a
25 potato SBE gene. The 5' region from nucleotides 1 to 2082 contain the promoter region of the SBE gene. A TATA box candidate at nucleotide 2048 to 2051 is boxed. The homology between a potato SBE cDNA clone (Poulsen & Kreiberg (1993) Plant Physiol 102: 1053-1054) and the exon DNAs begin at 2083 bp and end at 9666 bp. The homology between the cDNA and the exon DNA is indicated by nucleotides in
30 upper case letters, while the translated amino acid sequences are shown in the single letter code below the exon DNA. Intron sequences are indicated by lower case letters.

Figure 7 is a plasmid map of pBEA7, which is 9.54 k base pairs in size. Plasmid pBEA 11 comprises the first intron sequence of the potato SBE gene. This first intron sequence, which has 1177 base pairs, is shown in Figure 3 and lies between the first exon and the second exon.

These experiments and aspects of the present invention are now discussed in more detail.

EXPERIMENTAL PROTOCOL

ISOLATION, SUBCLONING IN PLASMIDS, AND SEQUENCING OF GENOMIC SBE CLONES

Various clones containing the potato SBE gene were isolated from a Desiree potato genomic library (Clontech Laboratories Inc., Palo Alto CA, USA) using radioactively labelled potato SBE cDNA (Poulsen & Kreiberg (1993) Plant Physiol. 102:1053-1054) as probe. The fragments of the isolated λ -phages containing SBE DNA (λ SBE 3.2 - NCIMB 40751 - and λ SBE-3.4 - NCIMB 40752) were identified by Southern analysis and then subcloned into pBluescript II vectors (Clontech Laboratories Inc., Palo Alto CA, USA). λ SBE 3.2 contains a 15 kb potato DNA insert and λ SBE-3.4 contains a 13 kb potato DNA insert. The resultant plasmids were called pGB3, pGB11, pGB15, pGB16 and pGB25 (see discussion below). The respective inserts were then sequenced using the Pharmacia Autoread Sequencing Kit (Pharmacia, Uppsala) and a A.L.F. DNA sequencer (Pharmacia, Uppsala).

In total, a stretch of 11.5 kb of the SBE gene was sequenced. The sequence was deduced from the above-mentioned plasmids, wherein: pGB25 contains the sequences from 1 bp to 836 bp, pGB15 contains the sequences from 735 bp to 2580 bp, pGB16 contains the sequences from 2580 bp to 5093 bp, pGB11 contains the sequences from 3348 bp to 7975 bp, and pGB3 contains the sequences from 7533 bp to 11468 bp.

In more detail, pGB3 was constructed by insertion of a 4 kb *EcoRI* fragment isolated from λ SBE 3.2 into the *EcoRI* site of pBluescript II SK (+). pGB11 was constructed by insertion of a 4.7 kb *XhoI* fragment isolated from λ SBE 3.4 into the *XhoI* site of pBluescript II SK (+). pGB15 was constructed by insertion of a 1.7 kb *SpeI* fragment isolated from λ SBE 3.4 into the *SpeI* site of pBluescript II SK (+). pGB16 was constructed by insertion of a 2.5 kb *SpeI* fragment isolated from λ SBE 3.4 into the *SpeI* site of pBluescript II SK (+). For the construction of pGB25 a PCR fragment was produced with the primers

5' GGA ATT CCA GTC GCA GTC TAC ATT AC 3'

and

5' CGG GAT CCA GAG GCA TTA AGA TTT CTG G 3'

and λ SBE 3.4 as a template.

The PCR fragment was digested with *BamHI* and *EcoRI*, and inserted in pBluescript II SK (+) digested with the same restriction enzymes.

CONSTRUCTION OF PLASMID pBEA11

The SBE intron 1 was amplified by PCR using the oligonucleotides

5' CGG GAT CCA AAG AAA TTC TCG AGG TTA CAT GG 3'

and

5' CGG GAT CCG GGG TAA TTT TTA CTA ATT TCA TG 3'

and the λ SBE 3.4 phage containing the SBE gene as template.

The PCR product was digested with *Bam*HI and inserted in a sense orientation in the *Bam*HI site of plasmid pPATA1 (described in WO 94/24292) between the patatin promoter and the 35S terminator. This construction, pABE7, was digested with *Kpn*I, and the 2.4 kb "patatin promoter-SBE intron 1- 35S terminator" *Kpn*I fragment was isolated and inserted in the *Kpn*I site of the plant transformation vector pVictorIV Man yielding plasmid pBEA11.

PRODUCTION OF TRANSGENIC POTATO PLANTS

Axenic stock cultures

Shoot cultures of *Solanum tuberosum* 'Bintje' and 'Dianella' are maintained on a substrate (LS) of a formula according to Linsmaier, E.U. and Skoog, F. (1965), *Physiol. Plant.* 18: 100-127, in addition containing 2 μ M silver thiosulphate at 25°C and 16 h light/8 h dark.

The cultures were subcultured after approximately 40 days. Leaves were then cut off the shoots and cut into nodal segments (approximately 0.8 cm) each containing one node.

Inoculation of potato tissues

Shoots from approximately 40 days old shoot cultures (height approximately 5-6 cms) were cut into internodal segments (approximately 0.8 cm). The segments were placed into liquid LS-substrate containing the transformed *Agrobacterium tumefaciens* containing the binary vector of interest. The *Agrobacterium* were grown overnight in YMB-substrate (di-potassium hydrogen phosphate, trihydrate (0.66 g/l); magnesium sulphate, heptahydrate (0.20 g/l); sodium chloride (0.10 g/l); mannitol (10.0 g/l); and yeast extract (0.40 g/l)) containing appropriate antibiotics (corresponding to the resistance gene of the *Agrobacterium* strain) to an optical density at 660 nm (OD-660) of approximately 0.8, centrifuged and resuspended in the LS-substrate to an OD-660 of 0.5.

The segments were left in the suspension of *Agrobacterium* for 30 minutes and then the excess of bacteria were removed by blotting the segments on sterile filter paper.

Co-cultivation

5

The shoot segments were co-cultured with bacteria for 48 hours directly on LS-substrate containing agar (8.0 g/l), 2,4-dichlorophenoxyacetic acid (2.0 mg/l) and trans-zeatin (0.5 mg/l). The substrate and also the explants were covered with sterile filter papers, and the petri dishes were placed at 25°C and 16 h light/ 8 dark.

10

"Washing" procedure

After the 48 h on the co-cultivation substrate the segments were transferred to containers containing liquid LS-substrate containing 800 mg/l carbenicillin. The containers were gently shaken and by this procedure the major part of the *Agrobacterium* was either washed off the segments and/or killed.

15

Selection

After the washing procedure the segments were transferred to plates containing the LS-substrate, agar (8 g/l), trans-zeatin (1-5 mg/l), gibberellic acid (0.1 mg/l), carbenicillin (800 mg/l), and kanamycin sulphate (50-100 mg/l) or phosphinotricin (1-5 mg/l) or mannose (5 g/l) depending on the vector construction used. The segments were sub-cultured to fresh substrate each 3-4 weeks. In 3 to 4 weeks, shoots develop from the segments and the formation of new shoots continued for 3-4 months.

20
25

Rooting of regenerated shoots

The regenerated shoots were transferred to rooting substrate composed of LS-substrate, agar (8 g/l) and carbenicillin (800 mg/l).

30

The transgenic genotype of the regenerated shoot were verified by testing the rooting ability on the above mentioned substrates containing kanamycin sulphate (200 mg/l), by performing NPTII assays (Radke, S. E. et al, Theor. Appl. Genet. (1988), 75: 685-694) or by performing PCR analysis according to Wang *et al* (1993, NAR 21 pp 4153-4154). Plants which were not positive in any of these assays were discarded or used as controls. Alternatively, the transgenic plants could be verified by performing a GUS assay on the co-introduced β -glucuronidase gene according to Hodal, L. *et al*. (Pl. Sci. (1992), 87: 115-122).

10 Transfer to soil

The newly rooted plants (height approx. 2-3 cms) were transplanted from rooting substrate to soil and placed in a growth chamber (21°C, 16 hour light 200-400uE/m²/sec). When the plants were well established they were transferred to the greenhouse, where they were grown until tubers had developed and the upper part of the plants were senescing.

Harvesting

20 The potatoes were harvested after about 3 months and then analysed.

BRANCHING ENZYME ANALYSIS

25 The SBE expression in the transgenic potato lines were measured using the SBE assays described by Blennow and Johansson (Phytochemistry (1991) 30:437-444) and by standard Western procedures using antibodies directed against potato SBE.

STARCH ANALYSIS

30 Starch was isolated from potato tubers and analysed for the amylose:amylopectin ratio (Hovenkamp-Hermelink et al. (1988) Potato Research 31:241-246). In addition, the chain length distribution of amylopectin was determined by analysis of isoamylase

digested starch on a Dionex HPAEC. The number of reducing ends in isoamylase digested starch was determined by the method described by N. Nelson (1944) J. Biol.Chem. 153:375-380.

5 The results revealed that there was a reduction in the level of synthesis of SBE and/or the level of activity of SBE and/or the composition of starch SBE in the transgenic plants.

CONSTRUCTION OF SBE PROMOTER CONSTRUCT

10

An SBE promoter fragment was amplified from λ -SBE 3.4 using primers:

5' CCA TCG ATA CTT TAA GTG ATT TGA TGG C 3'

15

and

5' CGG GAT CCT GTT CTG ATT CTT GAT TTC C 3'.

20

The PCR product was digested with *Cla*I and *Bam*HI. The resultant 1.2 kb fragment was then inserted in pVictor5a (see Figure 9) linearised with *Cla*I and *Bg*II yielding pBEP2 (see Figure 10).

STARCH BRANCHING ENZYME MEASUREMENTS OF POTATO TUBERS

25

Potatoes from potato plants transformed with pBEA11 were cut in small pieces and homogenised in extraction buffer (50 mM Tris-HCl pH 7.5, Sodium-dithionit (0.1 g/l), and 2 mM DTT) using a Ultra-Turax homogenizer; 1 g of Dowex xl. was added pr. 10 g of tuber. The crude homogenate was filtered through a miracloth filter and centrifuged at 4°C for 10 minutes at 24.700 g. The supernatant was used for starch branching enzyme assays.

30

The starch branching enzyme assays were carried out at 25 °C in a volume of 400 μ l composed of 0.1 M Na citrate buffer pH 7.0, 0.75 mg/ml amylose, 5 mg/ml bovine serum albumin and the potato extract. At 0, 15 30 and 60 minutes aliquots of 50 μ l were removed from the reaction into 20 μ l 3 N HCl. 1 ml of iodine solution
5 was added and the decrease in absorbance at 620 nm was measured with an ELISA spectrophotometer.

The starch branching enzyme (SBE) levels in tuber extracts were measured from 24 transgenic *Dianella* potato plants transformed with plasmid pBEA11.
10

The results showed that the BEA11 transgenic lines produced tubers which have SBE levels that are only 10 % to 15 % of the SBE levels found in non transformed *Dianella* plants.

15 SUMMATION

The above-mentioned examples relate to the isolation and sequencing of a gene for potato SBE. The examples further demonstrate that it is possible to prepare SBE intron constructs. These SBE intron constructs can be introduced into plants, such
20 as potato plants. After introduction, a reduction in the level of synthesis of SBE and/or the level of activity of SBE and/or the composition of starch in plants can be achieved.

Without wishing to be bound by theory it is believed that the expressed sense intron
25 nucleotide sequence according to the present invention affects enzymatic activity *via* co-suppression and/or trans-activation. Reviews of these mechanisms has been published by Finnegan and McElroy (1994 *Biotechnology* 12 pp 883 - 887) and Matzke and Matzke (1995 *TIG* 11 No. 1 pp 1 - 3). By these mechanisms, it is believed that the sense introns of the present invention reduce the level of plant
30 enzyme activity (in particular SBE activity), which in turn for SBE activity is believed to influence the amylose:amylopectin ratio and thus the branching pattern of amylopectin.

Thus, the present invention provides a method wherein it is possible to manipulate the starch composition in plants, or tissues or cells thereof, such as potato tubers, by reducing the level of SBE activity by using sense intron sequences.

5 In summation the present invention therefore relates to the surprising use of sense intron sequences in a method to affect enzymatic activity in plants.

Other modifications of the present invention will be apparent to those skilled in the art without departing from the scope of the present invention. For example, it may
10 be possible to use antisense promoter sequences to affect enzymatic activity, such as antisense SBE promoter - such as a nucleotide sequence comprising the nucleotide sequence shown as SEQ. I.D. No. 28 or a variant, derivative or homologue thereof.

The following pages present a number of sequence listings which have been
15 consecutively numbered from SEQ.I.D. No. 1 - SEQ.I.D. No. 29. In brief, SEQ.I.D. No. 1 - SEQ.I.D. No. 13 represent sense intron sequences (genomic DNA); SEQ.I.D. No. 14 represents the SBE promoter sequence (genomic sequence); SEQ.I.D. No. 15 - SEQ.I.D. No. 27 represent antisense intron sequences; and SEQ.
I.D. No. 28 represents the sequence complementary to the SBE promoter sequence -
20 i.e. the SBE promoter sequence in antisense orientation. The full genomic nucleotide sequence for SBE including the promoter, exons and introns is shown as SEQ. I.D. No. 29 (see Figures 3 and 8 which highlight particular gene features).

SEQUENCE INFORMATION

SEQ.I.D. No. 1

Intron 1 sequence (1167 bp).

GTAATTTTACTAATTTTCATGTTAATTTCAATTATTTTTCAGCCTTTGCATTTTCATTTTCCAATATATCT
GGATCATCTCCTTAGTTTTTTTATTTTATTTTATAATATCAAATATGGAAGAAAAATGACACTTGTAG
AGCCATATGTAAGTATCATGTGACAAATTTGCAAGGTGGTTGAGTGTATAAAATTCAAAAATTGAGAGA
TGGAGGGGGGTGGGGGBARAGACAATATTTAGAAAGAGTGTCTAGGAGGTTATGGAGGACACGGATG
AGGGGTAGAAGGTTAGTTAGGTATTTGAGTGTGTCTGGCTTATCCTTTCATACTAGTAGTCGTGGAAT
TATTTGGGTAGTTTCTTGTTTTGTTATTTGATCTTTGTTATTCTATTTTCTGTTTCTTGTAATTCGATT
ATTGTATTATATATCTTGTCTGTTAGTTATTGTTCTCGGTAAGAATGCTCTAGCATGCTTCCTTTAGTGT
TTTATCATGCCTTCTTTATATTGCGTGTGCTTTGAAATGCTTTACTTTAGCCGAGGGTCTATTAGAAA
CAATCTCTCTATCTCGTAAGGTAGGGGTAAAGTCTCCACCACACTCCACTTGTGGGATTACATTGTGTT
TGTTGTTGTAAATCAATTATGTATACATAATAAGTGGATTTTTTACAACACAAATACATGGTCAAGGGC
AAAGTTCGAACACATAAAGGGTTCATTATATGTCCAGGGATATGATAAAAATTGTTTCTTTGTGAAAG
TTATATAAGATTTGTTATGGCTTTTGCTGGAAACATAATAAGTTATAATGCTGAGATAGCTACTGAAGT
TTGTTTTTTCTAGCCTTTTAAATGTACCAATAATAGATTCCGTATCGAACGAGTATGTTTTGATTACCT
GGTCATGATGTTTCTATTTTTTACATTTTTTGGTGTGAACTGCAATTGAAAATGTTGTATCCTATGA
GACGGATAGTTGAGAATGTGTTCTTTGTATGGACCTTGAGAAGCTCAAACGCTACTCCAATAATTTCTA
TGAATTCAAATTCAGTTTATGGCTACCAGTCAGTCCAGAAATTAGGATATGCTGCATATACTTGTTCAA
TTATACTGTAAATTTCTTAAGTTCTCAAGATATCCATGTAACCTCGAGAATTTCTTTGACAG

SEQ.I.D. No. 2

Intron 2 sequence (321 bp).

GTATGTTTGATAAATTTATATGGTTGCATGGATAGTATATAAATAGTTGGAAAACCTTCTGGACTGGTGCT
CATGGCATATTTGATCTGTGCACCGTGTGGAGATGTCAAACATGTGTTACTTCGTTCCGCCAATTTATA
ATACCTTAACTTGGGAAAGACAGCTCTTACTCCTGTGGGCATTTGTTATTTGAATTACAATCTTTATG
AGCATGGTGTTTTACATTATCAACTTCTTTCATGTGGTATATAACAGTTTTTAGCTCCGTTAATACCT
TTCTTCTTTTGTATATAAACTAACTGTGGTGCATTGCTTGCBBKK

SEQ.I.D. No. 3

Intron 3 sequence (504 bp).

GTAACAGCCAAAAGTTGTGCTTTAGGCAGTTTGACCTTATTTTGAAGATGAATTGTTTATACCTACTT
TGACTTTGCTAGAGAATTTTGCATACCGGGGAGTAAGTAGTGGCTCCATTTAGGTGGCACCTGGCCATT
TTTTTGATCTTTTAAAAGCTGTTTGATTGGGTCTTCAAAAAGTAGACAAGGTTTTTGGAGAAGTGAC
ACACCCCCGGAGTGTGAGTGGCAAAGCAAAGATTTTCACTAAGGAGATTCAAAATATAAAAAAGTATA
GACATAAAGAAGCTGAGGGGATTCAACATGTACTATACAAGCATCAAATATAGTCTTAAAGCAATTTTG
TAGAAATAAAGAAAGTCTTCTTCTGTTGCTTCACAATTTCTTCTATTATCATGAGTTACTCTTTCTG
TTCGAAATAGCTTCCTTAATATTAATTCATGATACTTTTGTGAGATTTAGCAGTTTTTCTTGTGTA
AACTGCTCTCTTTTTTTGCAG

SEQ.I.D. No. 4

Intron 4 sequence (146 bp).

GTAGGTCCTCGTCTACTACAAAATAGTAGTTTCCATCATCATAACAGATTTTCCTATTAAAGCATGATG
TTGCAGCATCATTGGCTTTCTTACATGTTCTAATTGCTATTAAGGTTATGCTTCTAATTAACATCCCA
CAATGCAG

SEQ.I.D. No. 5

Intron 5 sequence (218 bp).

GTTTTGTTATTTCATACCTTGAAGCTGAATTTTGAACACCATCATCACAGGCATTTTCGATTCATGTTCTT
ACTAGTCTTGTTATGTAAGACATTTTGAATGCAAAAGTTAAAATAATTGTGTCTTTACTAATTTGGAC
TTGATCCCATACTCTTCCCTTAACAAAATGAGTCAATTCTATAAGTGCTTGAGAACTTACTACTTCAG
CAATTAAACAG

SEQ.I.D. No. 6

Intron 6 sequence (198 bp).

GTATTTTAAATTTATTTCTACAATAAATTCTCAGAACAATTGTTAGATAGAATCCAAATATATAC
GTCCTGAAAGTATAAAAGTACTTATTTTCGCCATGGGCCTTCAGAATATTGGTAGCCGCTGAATATCAT
GATAAGTTATTTATCCAGTGACATTTTATGTTCACTCCTATTATGTCTGCTGGATACAG

SEQ.I.D. No. 7

Intron 7 sequence (208bp)

GTTTGTCTGTTTCTATTGCATTTTAAGGTTTCATATAGGTTAGCCACGGAAAATCTCACTCTTTGTGAGG
TAACCAGGGTCTGATGGATTATTCAATTTTCTCGTTTATCATTTGTTTATTCTTTTCATGCATTGTGT
TTCTTTTCAATATCCCTCTTATTTGGAGGTAATTTTCTCATCTATTCACTTTTAGCTTCTAACCACAG

SEQ.I.D. No. 8

Intron 8 sequence (293 bp).

GTATGTCTTACATCTTTAGATATTTTGTGATAATTACAATTAGTTTGGCTTACTTGAACAAGATTCATT
CCTCAAAATGACCTGAACTGTTGAACATCAAAGGGGTTGAAACATAGAGGAAAACAACATGATGAATGT
TTCCATTGTCTAGGGATTTCTATTATGTTGCTGAGAACAAATGTCATCTTAAAAAAACATTGTTTACT
TTTTTGTAGTATAGAAGATTACTGTATAGAGTTTGCAAGTGTGTCTGTTTTGGAGTAATTGTGAAATGT
TTGATGAACTTGATACAG

SEQ.I.D. No. 9

Intron 9 sequence (376 bp).

GTTCAAGTATTTTGAATCGCAGCTTGTTAAATAATCTAGTAATTTTATAGATTGCTTACTTGGAAGTCTA
CTTGGTCTGGGGATGATAGCTCATTTTCATCTTGTTCTACTTATTTTCCAACCGAATTTCTGATTTTGG
TTTCGAGATCCAAGTATTAGATTCATTTACACTTATTACCGCCTCATTTCTACCACTAAGGCCTTGATG
AGCAGCTTAAGTTGATTCTTTGAAGCTATAGTTTCAGGCTACCAATCCACAGCCTGCTATATTTGTTGG

ATACTTACCTTTTCTTTACAATGAAGTGATACTAATTGAAATGGTCTAAATCTGATATCTATATTTCTC
CGTCTTTCCTCCCCCTCATGATGAAATGCAG

SEQ.I.D. No. 10

Intron 10 sequence (172 bp).

GTAAATCATCTAAAGTTGAAAGTGTGGGTTTATGAAGTGCTTTAATTCTATCCAAGGACAAGTAGAA
ACCTTTTACCTTCCATTTCTTGATGATGGATTTTCATATTATTTAATCCAATAGCTGGTCAAATTCGGT
AATAGCTGTACTGATTAGTTACTTCACTTTGCAG

SEQ.I.D. No. 11

Intron 11 sequence (145 bp).

GTATATATGTTTTACTTATCCATGAAATTATTGCTCTGCTTGTTTTAATGTACTGAACAAGTTTTATG
GAGAAGTAACTGAAACAAATCATTTTACATTGTCTAATTTAACTCTTTTTTCTGATCCTCGCATGACG
AAACAG

SEQ.I.D. No. 12

Intron 12 sequence (242 bp).

GTAAGGATTTGCTTGAATAACTTTTGATAATAAGATAACAGATGTAGGGTACAGTTCTCTCACCAAAA
GAAGTGTAATTGTCTCATCCATCTTTAGTTGTATAAGATATCCGACTGTCTGAGTTCGGAAGTGTTTGA
GCCTCCTGCCCTCCCCCTGCGTTGTTTAGCTAATTCAAAAAGGAGAAAAGTGTATTATGATGATCTTTG
TCTTCATGCTGACATACAATCTGTTCTCATGACAG

SEQ.I.D. No. 13

Intron 13 sequence (797 bp).

GTACAGTTCTTGCCGTGTGACCTCCCTTTTTATTGTGGTTTTGTTTCATAGTTATTTGAATGCGATAGAA
GTTAACTATTGATTACCGCCACAATCGCCAGTTAAGTCTCTGAACTACTAATTTGAAAGGTAGGAATA
GCCGTAATAAGGTCTACTTTTGGCATCTTACTGTTACAAAACAAAAGGATGCCAAAAAATTTCTTCTCT
ATCCTCTTTTTCCCTAAACAGTGCATGTAGCTTGCACCTGCATAAACTTAGGTAAATGATCAAAAATG
AAGTTGATGGGAACTTAAACCGCCCTGAAGTAAAGCTAGGAATAGTCATATAATGTCCACCTTTGGTG
TCTGCGCTAACATCAACAACAACATACCTCGTGTAGTCCCAAAAGTGGTTTCAGGGGGAGGGTAGAGT
GTATGCAAAACTTACTCCTATCTCAGAGGTAGAGAGGATTTTTTCAATAGACCCTTGGCTCAAGAAAA
AAGTCCAAAAAGAAGTAACAGAAGTGAAGCAACATGTGTAGCTAAAGCGACCAACTTGTTTGGGACT
GAAGTAGTTGTTGTTGTTGAACAGTGCATGTAGATGAACACATGTCAGAAAATGGACAACACAGTTAT
TTTGTGCAAGTCAAAAAAATGTACTACTATTTCTTTGTGCAGCTTTATGTATAGAAAAGTTAAATACT
AATGAATTTTGCTAGCAGAAAAATAGCTTGGAGAGAAATTTTTATATTGAACCTAAGCTAATATATTC
ATCTTCTTTTTGCTTCTTCTCTCCTTGTGTTGTGAAG

SEQ.I.D. No. 14

DNA sequence of the SBE gene promoter region.

```
ATCATGGCCAATTACTGGTTCAAATGCATTACTTCCTTTTCAGATTCTTTCGAGTTCTCAT      60
GACCGGTCCTACTACAGACGATACTAACCGTGGAAGTGTTCATCTGCTTCTTAGAACT      120
CTATGGCTATTTTCGTTAGCTTGGCGTCGGTTTGAACATAGTTTTTGTTCCTTCAAACCTCTT      180
CATTTACAGTCAAAATGTTGTATGGTTTTTGTTCCTCAATGATGTTTACAGTGTGTG      240
TTGTCATCTGTACTTTTGCCTATTACTTGTTCGAGTTACATGTTAAAAAAGTGTATT      300
TTGCCATATTTTGTCTCTTATTATTATTATCATACATACATTATTACAAGGAAAAGACA      360
AGTACACAGATCTTAACGTTTATGTTCAATCAACTTTTGGAGGCATTGACAGGTACCACA      420
AATTTTGAGTTTATGATTAAGTTCAATCTTAGAATATGAATTTAACATCTATTATAGATG      480
CATAAAAATAGCTAATGATAGAACATTGACATTTGGCAGAGCTTAGGGTATGGTATATCC      540
AACGTTAATTTAGTAATTTTGTACGTACGTATATGAAATATTGAATTAATCACATGAA      600
CGGTGGATATTATATTATGAGTTGGCATCAGCAAAATCATTGGTGTAGTTGACTGTAGTT      660
GCAGATTTAATAATAAAATGGTAATTAACGGTCGATATTAATAAATACTCTCATTTCAAGT      720
GGGATTAGAACTAGTTATTAATAAAATGTATACTTTAAGTGATTTGATGGCATATAATTT      780
AAAGTTTTTCATTCATGCTAAAATTGTTAATTATTGTAATGTAGACTGCGACTGGAATT      840
ATTATAGTGTAATTTATGCATTCAAGTGTAATAATTAAGTATTGAACCTGTCTGTTTTAG      900
AAAATACTTTTACTTTAATATAGGATTTTGTTCATGCGAATTTAAATTAATCGATATTGA      960
ACACGGAATACCAAAATTAATAAGGATACACATGGCCTTCATATGAACCGTGAACCTTTG      1020
ATAACGTGGAAGTTCAAAGAAGGTAAAGTTTAAAGAATAAACTGACAAATTAATTTCTTTT      1080
ATTTGGCCCACTACTAAATTTGCTTTACTTTCTAACATGTCAAGTTGTGCCCTCTTAGTT      1140
GAATGATATTTCATTTTCATCCATAAGTTCAATTTGATTGTACATACCACCCATGATGTT      1200
CTGAAAAATGCTTGGCCATTCACAAAGTTTATCTTAGTTCCCTATGAACCTTTATAAGAAGC      1260
TTTAATTTGACATGTTATTTATATTAGATGATATAATCCATGACCCAATAGACAAGTGTA      1320
TTAATATTGTAACCTTGTAATTGAGTGTGTCTACATCTTATTCAATCATTTAAGGTCATT      1380
AAAATAAATTATTTTTTGACATTCTAAAACCTTTAAGCAGAATAAATAGTTTATCAATTAT      1440
TAAAAACAAAAACGACTTATTTATAAATCAACAAACAATTTTAGATTGCTCCAACATAT      1500
TTTTCCAAATTAATGCAGAAAATGCATAATTTTATACTTGATCTTTATAGCTTATTTTT      1560
TTTAGCCTAACCAACGAATATTTGTAAACTCACAACCTTGATTAAAAGGGATTTACAACAA      1620
GATATATATAAGTAGTGACAAATCTTGATTTTAAATATTTAATTTGGAGGTCAAAATTT      1680
TACCATAATCATTTGTATTTATAATTAAATTTTAAATATCTTATTTATACATATCTAGTA      1740
AACTTTTAAATATACGTATATACAAAATATAAAATTATTGGCGTTCATATTAGGTCAATA      1800
AATCCTTAACCTATATCTGCCTTACCCTAGGAGAAAGTAAAAAACTCTTTACCAAAAATA      1860
CATGTATTATGTATACAAAAGTCGATTAGATTACCTAAATAGAAATTGTATAACGAGTA      1920
AGTAAGTAGAAATATAAAAAAACTACAATACTAAAAAAATATGTTTTACTTCAATTTTCG      1980
AAACTAATGGGGTCTGAGTGAAATATTCAGAAAGGGAGGACTAACAAAAGGTCATAAT      2040
GTTTTTTTATAAAAAGCCACTAAAATGAGGAAATCAAGAATCAGAACATACAAGAAGGCA      2100
GCAGCTGAAGCAAAGTACCATAATTTAATCAATGGAATTAATTTCAAAGTTTTATCAA      2160
ACCCATTCG
```

SEQ.I.D. No. 15

Intron 1 antisense sequence (1167 bp).

CTGTCAAAGAAATTCTCGAGGTTACATGGATATCTTGAGAACTTAAGAAATTTTACAGTATAATTGAAC
AAGTATATGCAGCATATCCTAATTTCTGGACTGACTGGTAGCCATAAACTGAATTTGAATTCATAGAAA
TTATTGGAGTAGCGTTTGAGCTTCTCAAGGTCCATACAAAGAACACATTCTCAACTATCCGTCTCATAG
GATACAACATTTTCAATTGCAGTTCAACACCAAAAAAATGTAAAAATAGAAACATCATGACCAGGTAA
TCAAAACATACTCGTTTCGATACGGAATCTATTATTGGTACATTTAAAAGGCTAGAAAAACAACTTCA
GTAGCTATCTCAGCATTATAACTTATTATGTTTCCAGCAAAAGCCATAACAAATCTTATATAACTTTCA
CAAAGAAACAATTTTTATCATATCCCTGGACATATAATGAACCCCTTTATGTGTTTCAGAACTTTGCCCTT
GACCATGTATTTGTGTTGTAAAAATCCACTTATTATGTATACATAATTGATTTACAACAACAAACACA
ATGTAATCCCAAGTGGAGTGTGGTGAGGACTTTACCCCTACCTTACGAGATAGAGAGATTGTTTCTA
ATAGACCCCTCGGCTAAAGTAAAAGCATTTCAAAGCAACGCGAATATAAAGAAGGCATGATAAAACACTA
AAGGAAGCATGCTAGAGCATTTCTTACCGAGGAACAATAACTACGACAAGATATATAATACAATAATCGA
AGTACAAGAAACAGAAAATAGAATAACAAAGATCAAATAACAAAACAAGAACTACCCAAATAATTTCCA
CGACTACTAGTATGAAAGGATAAGCCAGACAACACTCAAATACCTAACTAACCTTCTACCCCTCATCCG
TGTCTCCATAACCTCCTAGAACACTCTTTCTAAATATTGTCTYTVCCCCACCCCCCTCCATCTCTC
AATTTTTGAATTTTATACACTCAACCACCTTGCAAATTTGTCACATGATACTTACATATGGCTCTACAA
GTGTCAATTTTTCTTCCATATTTGATATTATAAAAAATAAAATAAAAACTAAGGAGATGATCCAGATAT
ATTGGAATGAAATGCAAAGGCTAAAAATAATTGAAATTAACATGAAATTAGTAAAAATTAC

SEQ.I.D. No. 16

Intron 2 antisense sequence (321 bp).

MMVVGCAAGCAATGCACCACAGTTAGTTTATATCAAAAAGAAGAAAGGTATTAAACGGAGCTAAAACTG
TTATATACCACATGAAAGAAGTTGATAATGTGAAAACACCATGCTCATAAAGATTGTAATTCAAATAAC
AAATGCCCACAGGAGTAAAGAGCTGTCTTTCCCAAGTTAAGGTATTATAAATTGGCGGAACGAAGTAAC
ACATGTTTGACATCTCCACACGGTGCACAGATCAAATATGCCATGAGCACCAGTCCAGAAGTTTTCCAA
CTATTTATATACTATCCATGCAACCATATAAATTATCAAACATAC

SEQ.I.D. No. 17

Intron 3 antisense sequence (504 bp).

CTGCAAAAAAGAGAGCAGTTTACACAAGAAAAAAGTCTAAATCTCAACAAAAGTATCATGAATTTAA
TATTAAGGAAGCTATTTGGAACAGAAAGAGTAACTCATGATAATAGAAGGAAATTGTGAAGCAACAGAA
GGAAGACTTTCTTTATTTCTACAAAATTGCTTTAAGACTATATTTGATGCTTGTATAGTACATGTTGAA
TCCCTCAGCTTCTTTATGTCTATACTTTTTTATATTTGAATCTCCTTAGTGAAATCTTTGCTTTG
CCACTGACACTCCGGGGGTGTGTCACTTCTCAAAAACCTTGTCTACTTTTTTGAAGACCCAATCAAAC
AGCTTTTTTAAAGATCAAAAAAATGGCCAGGTGCCACCTAAATGGAGCCACTACTTACTCCCCGGTATG
CAAATTTCTTAGCAAAGTCAAAGTAGGTATAAACAATTCATCTTCCAAAATAAGGTCAAAGTGCCTAA
AGCACAACTTTTGGCTGTTAC

SEQ.I.D. No. 18

Intron 4 antisense sequence (146 bp).

CTGCATTGTGGATGAGTTAATTAGAAGCATAACCTTAATAGCAATTAGAACATGTAAGAAAGCCAATGA
TGCTGCAACATCATGCTTTAATAGGAAAATCTGTTATGATGATGGAACTACTATTTTGTAGTAGACCGA
GGACCTAC

SEQ.I.D. No. 19

Intron 5 antisense sequence (218 bp).

CTGTTTAATTGCTGAAGTAGTAAGTTCTCAAGCACTTATAGAATTGACTCATTTTGTAAAGGGAAAGAG
TATGGGATCAAGTCCAAATTAGTAAAGACACAATTATTTTAACTTTTGCATTTCAAATGTCTTACATA
ACAAGACTAGTAAGAACATGAATCGAAATGCCTGTGATGATGGTGTTCAAAATTCAGCTTCAAGGTATG
AATAACAAAAC

SEQ.I.D. No. 20

Intron 6 antisense sequence (198 bp).

CTGTATCCAGCAGACATAATAGGAGTGAACATAAAAAATGTCACTGGATAAAATAACTTATCATGATATTC
AGCGGCTACCAATATTCTGAAGGCCCATGGCGAAAATAAGTACTTTTATACTTTTCAGGACGTATATATT
TGGATTCTATCTAACAATTGTTCTGAGAATTATTTAGTTGTAGAAATAAATTTAAAATAC

SEQ.I.D. No. 21

Intron 7 antisense sequence (208 bp).

CTGTGGTTAGAAGCTAAAAGTGAATAGATGAGAAAAATTACCTCCAAATAAGAGGGATATTGAAAAAGA
AACACAATGCATGAAAAGAATAAACAATGATAAACGAGAAAAATTGAATAATCCATCAGAACCCTGGTT
ACCTCACAAAGAGTGAGATTTTCCGTGGCTAACCTATATGAACCTTAAAATGCAATAGAAACAGACAAAC

SEQ.I.D. No. 22

Intron 8 antisense sequence (293 bp).

CTGTACAAGTTCATCAAACATTTTACAATTACTCCAAAACAGACACACTTGCAAACCTCTATACAGTAAT
CTTCTATACTACAAAAAAGTAAACAATGTTTTTTTAAAGATGACATTTGTTCTCAGCAACATAATAGAA
ATCCCTAGACAATGGAAACATTCATCATGTTGTTTTTCTCTATGTTTCAACCCCTTTGATGTTCAACAG
TTCAGGTCATTTTGAGGAATGAATCTTGTTCAAGTAAGCCAACTAATTGTAATTATCACAAAATATCT
AAAGATGTAAGACATAC

SEQ.I.D. No. 23

Intron 9 antisense sequence (376 bp).

CTGCATTTTCATCATGAGGGGGAGGAAAGACGGAGAAATATAGATATCAGATTTAGACCATTTCAATTAG
TATCACTTCATTGTAAAGAAAAGGTAAGTATCCAACAAATATAGCAGGCTGTGGATTGGTAGCCTGAAA
CTATAGCTTCAAAGAATCAACTTAAGCTGCTCATCAAGGCCTTAGTGGTAGAAATGAGGCGGTAATAAG
TGTAATGAATCTAATACTTGGATCTCGAAACAAAATCAGAAATTCGGTTGGAAAATAAGTAGAACAA

GATGAAATGAGCTATCATCCCCAGAACCAAGTAGACTTCCAAGTAAGCAATCTAAAAATTACTAGATTA
TTTAACAAGCTGCGATTCAAATACTTGAAC

SEQ.I.D. No. 24

Intron 10 antisense sequence (172 bp).

CTGCAAAGTGAAGTAACTAATCAGTACAGCTATTACCGAATTTGACCAGCTATTGGATTAAATAATATG
AAATCCATCATCAAGAAATGGAAGGTAAAAAGGTTTCTACTTGTCTTGGATAGAATTAAAGCACTTCA
TAAACCAACACTTTCAACTTTAGATGATTTTAC

SEQ.I.D. No. 25

Intron 11 antisense sequence (145 bp).

CTGTTTTCGTCATGCGAGGATCAGAAAAAGAGTTAAATTAGACAATGTGAAAATGATTGTTTCAGTT
ACTTCTCCATAAACTTGTTTCAGTACATTAAAAACAAGCAGAGCAATAATTTTCATGGATAAGTAAACA
TATATAC

SEQ.I.D. No. 26

Intron 12 antisense sequence (242 bp).

CTGTCATGAGAACAGATTGTATGTCAGCATGAAGACAAAGATCATCAATAAACAGTTTTCTCCTTTTTG
AATTAGCTAAACAACGCAGGGGGAGGGCAGGAGGCTCAAACACTTCCGAACTCAGACAGTCGGATATCT
TATACAACTAAAGATGGATGAGACAATTACAGTTCTTTTTGGTGAGAGAACTGTACCTACATCTGTTA
TCTTATTATCAAAAGTTATTCAAGCAAATCCTTAC

SEQ.I.D. No. 27

Intron 13 antisense sequence (797 bp).

CTTCACAAACAAGGAGAAGAAGCAAAAAGAAAGATGAATATAGTTAGCTTAGTTCAATATAAAAAA
TTTCTCTCCAAGCTATTTTTCTGCTAGCAAAATTCATTAGTTATTTAACTTTTCTATACATAAAGCTGC
ACAAAGAAATAGTAGTACATTTTTTTGACTTGACAAAATAACTGTGTTGTCCATTTTCTGACATGTGT
TCATCTACATGCACTGTTTCAACAACAACAACACTTTCAGTCCCAAACAAGTTGGGTCGCTTTAGCTAC
ACATGTTGCTTTCACTTCTGTTACTTCTTTTTGGACTTTTTTCTTGAGCCAAGGGTCTATTGAAAAAA
TCCTCTCTACCTCTGAGATAGGAGTAAGTTTTGCATACACTCTACCTCCCCCTGAAACCACTTTGTGG
GACTACACGAGGTATGTTGTTGTTGATGTTAGCGCAGACACCAAAGGTGGACATTATATGACTATTCCCT
AGCTTTACTTCAGGGCGGTTTTAAGTTCCCATCAACTTCATTTTGTATCATTTACCTAAGTTTATGCAG
GTGCAAGCTACATGCACTGGTTTAGGGAAAAAGAGGATAGAGAAGAATTTTTTGGCATCCTTTTGT
TGTAACAGTAAGATGCCAAAGTAGACCTTATTACGGCTATTCCTACCTTTCAAATTAGTAGTTCAGAG
GACTTAACTGGCGATTGTGGCGGTAATCAATAGTTAACTTCTATCGCATTCAAATAACTATGAACAAAA
CCACAATAAAAAGGGAGGTCACACGGCAAGAACTGTAC

SEQ.I.D. No. 28

Antisense DNA sequence of the SBE gene promoter region.

CGAATGGGTTTTGATAAACTTTGAAATTAATTTCCATTGATTAAATTATGGTACTTTGC	60
TTCAGCTGCTGCCTTCTTGTATGTTCTGATTCTTGATTTCCCTCATTTTAGTGGCTTTTTA	120
TAAAAAACATTATGACCCTTTTGTAGTCTCCCTTTCTGAATATTTCACTCAGACCC	180
CATTAGTTTCGAAATTGAAGTAAAACATATTTTTTTTAGTATTGTAGTTTTTTTATATTT	240
CTACTTACTTACTCGTTATACAATTTCTATTTAGGTAATCTAATCGACTTTTTGTATACA	300
TAATACATGTATTTTTGGTAAAGAGTTTTTACTTTCTCCTAGTGGTAAGGCAGATATAG	360
TTAAGGATTTATTGACCTAATATGAACGCCAATAATTTTATATTTGTATATACGTATAT	420
TTAAAAGTTTACTAGATATGTATAAATAAGATATTTAAAATTTAATTATAAATACAAATG	480
ATTATGGTAAAAATTTGACCTCCAAATTAAAATATTTAAAATCAAGATTTGTCACTACTT	540
ATATATATCTTGTGTAAATCCCTTTTAAATCAAGTTGTGAGTTTACAAATATTCGTTGGT	600
TAGGCTAAAAAAATAAGCTATAAAGATCAAGTATAAAATTATGCATTTTCTGCATTTAA	660
TTTGGAAAAATATGTTGGAGCAATCTAAAATGTGTTGTTGATTTATAAATAAGTCGTTTT	720
TTGTTTTTAATAATTGATAAACTATTTATTCTGCTTAAAGTTTTAGAATGTCAAAAAATA	780
ATTTATTTTAATGACCTTAAATGATTGAATAAGATGTAGACACACTCAATTACAAAGTTA	840
CAATATTAATACACTTGTCTATTGGGTCATGGATTATATCATCTAATATAAATAACATGT	900
CAAATTAAAGCTTCTTATAAAGTTCATAGGAATAAGATAAACTTTGTGAATGGCCAAGC	960
ATTTTTTCAGAACATCATGGGTGGTATGACAAATCAAATTGAACTTATGGGATGAAAAATGA	1020
ATATCATTTCAACTAAGAGGGCACAACCTTGACATGTTAGAAAGTAAAGCAAATTTAGTAGT	1080
GGGCCAAATAAAAGAAATTAATTTGTCTAGTTTATTCTTAACTTTACCTTCTTTGAACTT	1140
CCACGTTATCAAAGGTTACGGTTCATATGAAGGCCATGTGTATCCTTTTTAATTTTGGT	1200
ATTCGGTGTTCATATCGATTAAATTTAAATTCGCATGACAAAATCCTATATTAAAGTATA	1260
AAGTATTTTCTAAAACAGACAAGTTCAACTTTAATTTTACTGAATGCATAAATTTA	1320
CACTATAATAATTCCAGTCGCAGTCTACATTACAATAATTAACAATTTTAGCATGAAATG	1380
AAAACTTTAAATATATGCCATCAAATCACTTAAAGTATACATTTTTTTAATAACTAGT	1440
TCTAATCCCACTTGAAATGAGAGTTATTTTAATATCGACCGTTAATTACCATTTTATTAT	1500
TAAATCTGCAACTACAGTCAACTACACCAATGATTTTGCTGATGCCAACTCATAATATAA	1560
TATCCACCGTTCATGTGATTAATTCAATATTTTATATACGTACGTAACAAAAATTACTAA	1620
ATTAACGTTGGATATACCATAACCCTAAGCTCTGCCAAATGTCAATGTTCTATCATTAGCT	1680
ATTTTTATGCATCTATAATAGATGTTAAATTCATATTCTAAGATTGAACTTAATCATAAA	1740
CTCAAAATTTGTGGTACCTGTCAATGCCTCCAAAAGTTGATTGAACATAAACGTTAAGAT	1800
CTGTGTACTTGTCTTTTCCTTGTAATAATGTATGTATGATAATAATAAGAGAACAAA	1860
ATATGGCAAAATAAAACACTTTTTTAACATGTAACCTAAAACAAGTAATAGGCAAAAGTAC	1920
AGATGACAACACAACACTGTAAACATCATTTGAGGAAAACAAAAACCATAACAACATTTTGA	1980
CTGTAAATGAAGAGTTTGAAAACAAAACTATGTTCAAACCGACGCAAGCTAACGAAAA	2040
TAGCCATAGAGTTCTAAGAAGCAGATGCAACAGTTCCACGGGTTAGTATCGTCTGTAGTA	2100
GGACCGGTCTATGAGAACTCGAAAGAATCTGAAAGGAAGTAATGCATTTGAACCAGTAATT	2160
GGCCATGAT	

SEQ.I.D. No. 29

Genomic SBE gene

ATCATGGCCA	ATTACTGGTT	CAAATGCATT	ACTTCCTTTC	AGATTCTTTC	GAGTTCTCAT	60
GACCGGTCCT	ACTACAGACG	ATACTAACCC	GTGGAACGTG	TGCATCTGCT	TCTTAGAACT	120
CTATGGCTAT	TTTCGTTAGC	TTGGCGTCGG	TTTGAACATA	GTTTTTGTTT	TCAAACCTCTT	180
CATTTACAGT	CAAAATGTTG	TATGGTTTTT	GTTTTCTCTCA	ATGATGTTTA	CAGTGTGTGTG	240
TTGTCATCTG	TACTTTTGCC	TATTACTTGT	TTGAGTTAC	ATGTTAAAAA	AGTGTTTATT	300
TTGCCATATT	TTGTTCTCTT	ATTATTATTA	TCATACATAC	ATTATTACAA	GGAAAAGACA	360
AGTACACAGA	TCTTAACGTT	TATGTTCAAT	CAACTTTTGG	AGGCATTGAC	AGGTACCACA	420
AATTTTGAGT	TTATGATTAA	GTTCAATCTT	AGAATATGAA	TTTAACATCT	ATTATAGATG	480
CATAAAAATA	GCTAATGATA	GAACATTGAC	ATTTGGCAGA	GCTTAGGGTA	TGGTATATCC	540
AACGTTAATT	TAGTAATTTT	TGTTACGTAC	GTATATGAAA	TATTGAATTA	ATCACATGAA	600
CGGTGGATAT	TATATTATGA	GTTGGCATCA	GCAAAATCAT	TGGTGTAGTT	GACTGTAGTT	660
GCAGATTTAA	TAATAAAATG	GTAATTAACG	GTCGATATTA	AAATAACTCT	CATTTCAAGT	720
GGGATTAGAA	CTAGTTATTA	AAAAAATGTA	TACTTTAAGT	GATTTGATGG	CATATAATTT	780
AAAGTTTTTC	ATTTTCATGCT	AAAATTGTTA	ATTATTGTAA	TGTAGACTGC	GACTGGAATT	840
ATTATAGTGT	AAATTTATGC	ATTCAGTGTA	AAATTAAAGT	ATTGAACCTG	TCTGTTTTAG	900
AAAATACTTT	ATACTTTAAT	ATAGGATTTT	GTCATGCGAA	TTTAAATTAA	TCGATATTGA	960
ACACGGAATA	CCAAAATTAA	AAAGGATACA	CATGGCCTTC	ATATGAACCG	TGAACCTTTG	1020
ATAACGTGGA	AGTTCAAAGA	AGGTAAAGTT	TAAGAATAAA	CTGACAAATT	AATTTCTTTT	1080
ATTTGGCCCA	CTACTAAATT	TGCTTTACTT	TCTAACATGT	CAAGTTGTGC	CCTCTTAGTT	1140
GAATGATATT	CATTTTTTCAT	CCCATAAGTT	CAATTTGATT	GTCATACCAC	CCATGATGTT	1200
CTGAAAAATG	CTTGGCCATT	CACAAAGTTT	ATCTTAGTTC	CTATGAACTT	TATAAGAAGC	1260
TTTAATTTGA	CATGTTATTT	ATATTAGATG	ATATAATCCA	TGACCCAATA	GACAAGTGTA	1320
TTAATATTGT	AACTTTGTAA	TTGAGTGTGT	CTACATCTTA	TTCAATCATT	TAAGGTCATT	1380
AAAATAAATT	ATTTTTTGAC	ATTCTAAAAC	TTTAAGCAGA	ATAAATAGTT	TATCAATTAT	1440
TAAAAACAAA	AAACGACTTA	TTTATAAATC	AACAAACAAT	TTTAGATTGC	TCCAACATAT	1500
TTTTCCAAAT	TAAATGCAGA	AAATGCATAA	TTTTATACTT	GATCTTTATA	GCTTATTTTT	1560
TTTAGCCTAA	CCAACGAATA	TTTGTAACCT	CACAACTTGA	TTAAAAGGGA	TTTACAACAA	1620
GATATATATA	AGTAGTGACA	AATCTTGATT	TAAATATTTT	TAATTTGGAG	GTCAAAATTT	1680
TACCATAATC	ATTTGTATTT	ATAATTAAAT	TTTAAATATC	TTATTTATAC	ATATCTAGTA	1740
AACTTTTAAA	TATACGTATA	TACAAAATAT	AAAATTATTG	GCGTTCATAT	TAGGTCAATA	1800
AATCCTTAAC	TATATCTGCC	TTACCACTAG	GAGAAAGTAA	AAACTCTTTT	ACCAAAAATA	1860
CATGTATTAT	GTATACAAAA	AGTCGATTAG	ATTACCTAAA	TAGAAATTGT	ATAACGAGTA	1920
AGTAAGTAGA	AATATAAAAA	AACTACAATA	CTAAAAAATA	TATGTTTTAC	TTCAATTTTCG	1980
AAACTAATGG	GGTCTGAGTG	AAATATTCAG	AAAGGGGAGG	ACTAACAAAA	GGGTCATAAT	2040
GTTTTTTTAT	AAAAAGCCAC	TAAAATGAGG	AAATCAAGAA	TCAGAACATA	CAAGAAGGCA	2100
GCAGCTGAAG	CAAAGTACCA	TAATTTAATC	AATGGAAATT	AATTTCAAAG	TTTTATCAAA	2160
ACCCATTCGA	GGATCTTTTC	CATCTTTCTC	ACCTAAAGTT	TCTTCAGGGG	TAATTTTTTAC	2220
TAATTTTCATG	TTAATTTCAA	TTATTTTTAG	CCTTTGCATT	TCATTTTCCA	ATATATCTGG	2280
ATCATCTCCT	TAGTTTTTTA	TTTTATTTTT	TATAATATCA	AATATGGAAG	AAAAATGACA	2340
CTTGTTAGAGC	CATATGTAAG	TATCATGTGA	CAAATTTGCA	AGGTGGTTGA	GTGTATAAAA	2400
TTCAAAAATT	GAGAGATGGA	GGGGGGGTGG	GGGBARAGAC	AATATTTAGA	AAGAGTGTTC	2460
TAGGAGGTTA	TGGAGGACAC	GGATGAGGGG	TAGAAGGTTA	GTTAGGTATT	TGAGTGTGTG	2520

CTGGCTTATC	CTTTCATACT	AGTAGTCGTG	GAATTATTTG	GGTAGTTTCT	TGTTTTGTTA	2580
TTTGATCTTT	GTTATTCTAT	TTTCTGTTTC	TTGTACTTCG	ATTATTGTAT	TATATATCTT	2640
GTCGTAGTTA	TTGTTCCCTG	GTAAGAATGC	TCTAGCATGC	TTCCTTTAGT	GTTTTATCAT	2700
GCCTTCTTTA	TATTCGCGTT	GCTTTGAAAT	GCTTTTACTT	TAGCCGAGGG	TCTATTAGAA	2760
ACAACTCTCTC	TATCTCGTAA	GGTAGGGGTA	AAGTCCTCAC	CACACTCCAC	TTGTGGGATT	2820
ACATTGTGTT	TGTTGTTGTA	AATCAATTAT	GTATACATAA	TAAGTGGATT	TTTTACAACA	2880
CAAATACATG	GTCAAGGGCA	AAGTTCTGAA	CACATAAAGG	GTTCAATTATA	TGTCCAGGGA	2940
TATGATAAAA	ATTGTTTCTT	TGTGAAAGTT	ATATAAGATT	TGTTATGGCT	TTTGCTGGAA	3000
ACATAATAAG	TTATAATGCT	GAGATAGCTA	CTGAAGTTTG	TTTTTTCTAG	CCTTTTAAAT	3060
GTACCAATAA	TAGATTCCGT	ATCGAACGAG	TATGTTTTGA	TTACCTGGTC	ATGATGTTTC	3120
TATTTTTTAC	ATTTTTTTTG	TGTTGAACTG	CAATTGAAAA	TGTTGTATCC	TATGAGACGG	3180
ATAGTTGAGA	ATGTGTTCTT	TGTATGGACC	TTGAGAAGCT	CAAACGCTAC	TCCAATAATT	3240
TCTATGAATT	CAAATTCAGT	TTATGGCTAC	CAGTCAGTCC	AGAAATTAGG	ATATGCTGCA	3300
TATACTTGTT	CAATTATACT	GTAAAATTTT	TTAAGTTCTC	AAGATATCCA	TGTAACCTCG	3360
AGAATTTCTT	TGACAGGCTT	CTAGAAATAA	GATATGTTTT	CCTTCTCAAC	ATAGTACTGG	3420
ACTGAAGTTT	GGATCTCAGG	AACGGTCTTG	GGATATTTCT	TCCACCCCAA	AATCAAGAGT	3480
TAGAAAAGAT	GAAAGGGTAT	GTTTGATAAT	TTATATGGTT	GCATGGATAG	TATATAAATA	3540
GTTGAAAAAC	TTCTGGACTG	GTGCTCATGG	CATATTTGAT	CTGTGCACCG	TGTGGAGATG	3600
TCAAACATGT	GTTACTTCGT	TCCGCCAATT	TATAATACCT	TAACCTGGGA	AAGACAGCTC	3660
TTTACTCCTG	TGGGCATTTG	TTATTTGAAT	TACAATCTTT	ATGAGCATGG	TGTTTTCACA	3720
TTATCAACTT	CTTTCATGTG	GTATATAACA	GTTTTTAGCT	CCGTTAATAC	CTTCTTCTT	3780
TTTGATATAA	ACTAACTGTG	GTGCATTGCT	TGCBKKKATG	AAGCACAGTT	CAGCTATTTT	3840
CGCTGTTTTG	ACCGATGACG	ACAATTCGAC	AATGGCACCC	CTAGAGGAAG	ATGTCAAGAC	3900
TGAAAAATATT	GGCCTCCTAA	ATTTGGATCC	AACTTTGGAA	CCTTATCTAG	ATCACTTCAG	3960
ACACAGAATG	AAGAGATATG	TGGATCAGAA	AATGCTCATT	GAAAAATATG	AGGGACCCCT	4020
TGAGGAATTT	GCTCAAGGTA	ACAGCCAAAA	GTTGTGCTTT	AGGCAGTTTG	ACCTTATTTT	4080
GGAAGATGAA	TTGTTTATAC	CTACTTTGAC	TTTGCTAGAG	AATTTTGCAT	ACCGGGGAGT	4140
AAGTAGTGGC	TCCATTTAGG	TGGCACCTGG	CCATTTTTTT	GATCTTTTAA	AAAGCTGTTT	4200
GATTGGGTCT	TCAAAAAAGT	AGACAAGGTT	TTTGGAGAAG	TGACACACCC	CCGGAGTGTC	4260
AGTGGCAAAG	CAAAGATTTT	CACTAAGGAG	ATTCAAATAA	TAAAAAAGT	ATAGACATAA	4320
AGAAGCTGAG	GGGATTCAAC	ATGTACTATA	CAAGCATCAA	ATATAGTCTT	AAAGCAATTT	4380
TGTAGAAATA	AAGAAAGTCT	TCCTTCTGTT	GCTTCACAAT	TTCCCTTCTAT	TATCATGAGT	4440
TACTCTTTCT	GTTGAAATAA	GCTTCCTTAA	TATTAAATTC	ATGATACTTT	TGTTGAGATT	4500
TAGCAGTTTT	TTCTTGTTGA	AACTGCTCTC	TTTTTTTGCA	GGTTATTTAA	AATTTGGATT	4560
CAACAGGGAA	GATGGTTGCA	TAGTCTATCG	TGAATGGGCT	CCTGCTGCTC	AGTAGGTCCT	4620
CGTCTACTAC	AAAATAGTAG	TTCCATCAT	CATAACAGAT	TTTCCTATTA	AAGCATGATG	4680
TTGCAGCATC	ATTGGCTTTC	TTACATGTTT	TAATTGCTAT	TAAGGTTATG	CTTCTAATTA	4740
ACTCATCCAC	AATGCAGGGA	AGCAGAAGTT	ATTGGCGATT	TCAATGGATG	GAACGGTTCT	4800
AACCACATGA	TGGAGAAGGA	CCAGTTTGGT	GTTTGGAGTA	TTAGAATTCC	TGATGTTGAC	4860
AGTAAGCCAG	TCATTCCACA	CAACTCCAGA	GTTAAGTTTC	GTTTCAAACA	TGGTAATGGA	4920
GTGTGGGTAG	ATCGTATCCC	TGCTTGATAA	AAGTATGCCA	CTGCAGACGC	CACAAAGTTT	4980
GCAGCACCAT	ATGATGGTGT	CTACTGGGAC	CCACCACCTT	CAGAAAGGTT	TTGTTATTCA	5040
TACCTTGAAG	CTGAATTTTG	AACACCATCA	TCACAGGCAT	TTCGATTCAT	GTTCTTACTA	5100
GTCTTGTTAT	GTAAGACATT	TTGAAATGCA	AAAGTTAAAA	TAATTGTGTC	TTTACTAATT	5160
TGGACTTGAT	CCCATACTCT	TTCCCTTAAC	AAAATGAGTC	AATTCTATAA	GTGCTTGAGA	5220

ACTTACTACT	TCAGCAATTA	AACAGGTACC	ACTTCAAATA	CCCTCGCCCT	CCCAAACCCC	5280
GAGCCCCACG	AATCTATGAA	GCACATGTCG	GCATGAGCAG	CTCTGAGCCA	CGTGTAATTT	5340
CGTATCGTGA	GTTTGCAGAT	GATGTTTTAC	CTCGGATTAA	GGCAAATAAC	TATAATACTG	5400
TCCAGTTGAT	GGCCATAATG	GAACATTCTT	ACTATGGATC	ATTTGGATAT	CATGTTACAA	5460
ACTTTTTTGC	TGTGAGCAGT	AGATATGGAA	ACCCGGAGGA	CCTAAAGTAT	CTGATAGATA	5520
AAGCACATAG	CTTGGGTTTA	CAGGTTCTGG	TGGATGTAGT	TCACAGTCAT	GCAAGCAATA	5580
ATGTCACTGA	TGGCCTCAAT	GGCTTTGATA	TTGGCCAAGG	TTCTCAAGAA	TCCTACTTTC	5640
ATGCTGGAGA	GCGAGGGTAC	CATAAGTTGT	GGGATAGCAG	GCTGTTCAAC	TATGCCAATT	5700
GGGAGGTTCT	TCGTTTCCCT	CTTTCCAAC	TGAGGTGGTG	GCTAGAAGAG	TATAACTTTG	5760
ACGGATTTTCG	ATTTGATGGA	ATAACTTCTA	TGCTGTATGT	TCATCATGGA	ATCAATATGG	5820
GATTTACAGG	AACTATAAT	GAGTATTTCA	GCGAGGCTAC	AGATGTTGAT	GCTGTGGTCT	5880
ATTTAATGTT	GGCCAATAAT	CTGATTCA	AGATTTTCCC	AGATGCAACT	GTTATTGCCG	5940
AAGATGTTTC	TGGTATGCCG	GGCCTTGGCC	GGCCTGTTTC	TGAGGGAGGA	ATTGGTTTTC	6000
TTTACCGCCT	GGCAATGGCA	ATCCCAGATA	AGTGGATAGA	TTATTTAAAG	AATAAGAATG	6060
ATGAAGATTG	GTCCATGAAG	GAAGTAACAT	CGAGTTTGAC	AAATAGGAGA	TATACAGAGA	6120
AGTGATATAG	ATATGCGGAG	ACCCATGATC	AGGTATTTTA	AATTTATTTTC	TACAACTAAA	6180
TAATTCTCAG	AACAATTGTT	AGATAGAATC	CAAATATATA	CGTCCTGAAA	GTATAAAAGT	6240
ACTTATTTTC	GCCATGGGCC	TTCAGAATAT	TGGTAGCCGC	TGAATATCAT	GATAAGTTAT	6300
TTATCCAGTG	ACATTTTTAT	GTTCACTCCT	ATTATGTCTG	CTGGATACAG	TCTATTGTTG	6360
GTGACAAGAC	CATTGCATTT	CTCCTAATGG	ACAAAGAGAT	GTATTCTGGC	ATGTCTTGCT	6420
TGACAGATGC	TTCTCCTGTT	GTTGATCGAG	GAATTGCGCT	TCACAAGGTT	TGTCTGTTTC	6480
TATTGCATTT	TAAGGTTTCAT	ATAGGTTAGC	CACGGAAAT	CTCACTCTTT	GTGAGGTAAC	6540
CAGGGTTCTG	ATGGATTATT	CAATTTTCTC	GTTTATCATT	TGTTTATTCT	TTTCATGCAT	6600
TGTGTTTCTT	TTTCAATATC	CCTCTTATTT	GGAGGTAATT	TTTCTCATCT	ATTCATTTT	6660
AGCTTCTAAC	CACAGATGAT	CCATTTTTTC	ACAATGGCCT	TGGGAGGAGA	GGGGTACCTC	6720
AAATTCATGG	GTAACGAGGT	ATGTCTTACA	TCTTTAGATA	TTTTGTGATA	ATTACAATTA	6780
GTTTGGCTTA	CTTGAACAAG	ATTCATTCC	CAAAATGACC	TGAACTGTTG	AACATCAAAG	6840
GGGTGAAAC	ATAGAGGAAA	ACAACATGAT	GAATGTTTCC	ATTGTCTAGG	GATTTCTATT	6900
ATGTTGCTGA	GAACAAATGT	CATCTTAAAA	AAAACATTGT	TTACTTTTTT	GTAGTATAGA	6960
AGATTACTGT	ATAGAGTTTG	CAAGTGTGTC	TGTTTTGGAG	TAATTGTGAA	ATGTTTGATG	7020
AACTTGTACA	GTTTGGCCAT	CCTGAGTGGA	TTGACTTCCC	TAGAGAGGGC	AATAATTGGA	7080
GTTATGACAA	ATGTAGACGC	CAGTGGAACC	TCGCGGATAG	CGAACACTTG	AGATACAAGG	7140
TTCAAGTATT	TTGAATCGCA	GCTTGTTAAA	TAATCTAGTA	ATTTTATAGAT	TGCTTACTTG	7200
GAAGTCTACT	TGGTTCCTGGG	GATGATAGCT	CATTTTCATCT	TGTTCTACTT	ATTTTCCAAC	7260
CGAATTTCTG	ATTTTTGTTT	CGAGATCCAA	GTATTAGATT	CATTTTACACT	TATTACCGCC	7320
TCATTTCTAC	CACATAAGGCC	TTGATGAGCA	GCTTAAGTTG	ATTCTTTGAA	GCTATAGTTT	7380
CAGGCTACCA	ATCCACAGCC	TGCTATATTT	GTTGGATACT	TACCTTTTCT	TTACAATGAA	7440
GTGATACTAA	TTGAAATGGT	CTAAATCTGA	TATCTATATT	TCTCCGTCTT	TCCTCCCCCT	7500
CATGATGAAA	TGCAGTTTAT	GAATGCATTT	GATAGAGCTA	TGAATTCGCT	CGATGAAAAG	7560
TTCTCATTTCC	TCGCATCAGG	AAAACAGATA	GTAAGCAGCA	TGGATGATGA	TAATAAGGTA	7620
AAATCATCTA	AAGTTGAAAG	TGTTGGGTTT	ATGAAGTGCT	TTAATTCTAT	CCAAGGACAA	7680
GTAGAAACCT	TTTTACCTTC	CATTTCTTGA	TGATGGATTT	CATATTATTT	AATCCAATAG	7740
CTGGTCAAAT	TCGGTAATAG	CTGTACTGAT	TAGTTACTTC	ACTTTGCAGG	TTGTTGTGTT	7800
TGAACGTGGT	GACCTGGTAT	TTGTATTCAA	CTTCCACCCA	AAGAACACAT	ACGAAGGGTA	7860
TATATGTTTT	ACTTATCCAT	GAAATTATTG	CTCTGCTTGT	TTTTAATGTA	CTGAACAAGT	7920

TTTATGGAGA	AGTAACTGAA	ACAAATCATT	TTCACATTGT	CTAATTTAAC	TCTTTTTTCT	7980
GATCCTCGCA	TGACGAAAAC	AGGTATAAAG	TTGGATGTGA	CTTGCCAGGG	AAGTACAGAG	8040
TTGCACTGGA	CAGTGATGCT	TGGGAATTTG	GTGGCCATGG	AAGAGTAAGG	ATTTGCTTGA	8100
ATAACTTTTG	ATAATAAGAT	AACAGATGTA	GGGTACAGTT	CTCTCACCAA	AAAGAAGTGT	8160
AATTGTCTCA	TCCATCTTTA	GTTGTATAAG	ATATCCGACT	GTCTGAGTTC	GGAAGTGTTT	8220
GAGCCTCCTG	CCCTCCCCCT	GCGTTGTTTA	GCTAATTCAA	AAAGGAGAAA	ACTGTTTATT	8280
GATGATCTTT	GTCTTCATGC	TGACATACAA	TCTGTTCTCA	TGACAGACTG	GTCATGATGT	8340
TGACCATTTC	ACATCACCAG	AAGGAATACC	TGGAGTTCCA	GAAACAAATT	TCAATGGTCG	8400
TCCAAATTCC	TTCAAAGTGC	TGTCTCCTGC	GCGAACATGT	GTGGTACAGT	TCTTGCCGTG	8460
TGACCTCCCT	TTTTATTGTG	GTTTTGTTCA	TAGTTATTTG	AATGCGATAG	AAGTTAACTA	8520
TTGATTACCG	CCACAATCGC	CAGTTAAGTC	CTCTGAACTA	CTAATTTGAA	AGGTAGGAAT	8580
AGCCGTAATA	AGGTCTACTT	TTGGCATCTT	ACTGTTACAA	AACAAAAGGA	TGCCAAAAAA	8640
ATTCTTCTCT	ATCCTCTTTT	TCCCTAAACC	AGTGCATGTA	GCTTGCACCT	GCATAAACTT	8700
AGGTAAATGA	TCAAAAATGA	AGTTGATGGG	AACTTAAAAAC	CGCCCTGAAG	TAAAGCTAGG	8760
AATAGTCATA	TAATGTCCAC	CTTTGGTGTC	TGCGCTAACA	TCAACAACAA	CATACCTCGT	8820
GTAGTCCCAC	AAAGTGGTTT	CAGGGGGAGG	GTAGAGTGTA	TGCAAACTT	ACTCCTATCT	8880
CAGAGGTAGA	GAGGATTTTT	TCAATAGACC	CTTGGCTCAA	GAAAAAAAGT	CCAAAAAGAA	8940
GTAACAGAAG	TGAAAGCAAC	ATGTGTAGCT	AAAGCGACCC	AACTTGTTTG	GGACTGAAGT	9000
AGTTGTTGTT	GTTGAAACAG	TGCATGTAGA	TGAACACATG	TCAGAAAATG	GACAACACAG	9060
TTATTTTGTG	CAAGTCAAAA	AAATGTACTA	CTATTTCTTT	GTGCAGCTTT	ATGTATAGAA	9120
AAGTTAAATA	ACTAATGAAT	TTTGCTAGCA	GAAAAATAGC	TTGGAGAGAA	ATTTTTTATA	9180
TTGAACTAAG	CTAACTATAT	TCATCTTTCT	TTTTGCTTCT	TCTTCTCCTT	GTTTGTGAAG	9240
GCTTATTACA	GAGTTGATGA	ACGCATGTCA	GAAACTGAAG	ATTACCAGAC	AGACATTTGT	9300
AGTGAGCTAC	TACCAACAGC	CAATATCGAG	GAGAGTGACG	AGAACTTAA	AGATTCGTTA	9360
TCTACAAATA	TCAGTAACAT	TGACGAACGC	ATGTCAGAAA	CTGAAGTTTA	CCAGACAGAC	9420
ATTTCTAGTG	AGCTACTACC	AACAGCCAAT	ATTGAGGAGA	GTGACGAGAA	ACTTAAAGAT	9480
TCGTTATCTA	CAAATATCAG	TAACATTGAT	CAGACTGTTG	TAGTTTCTGT	TGAGGAGAGA	9540
GACAAGGAAC	TTAAAGATTC	ACCGTCTGTA	AGCATCATTA	GTGATGTTGT	TCCAGCTGAA	9600
TGGGATGATT	CAGATGCAAA	CGTCTGGGGT	GAGGACTAGT	CAGATGATTG	ATCGACCTT	9660
CTACCGATTG	GTGATCGCTA	TCCTTGCTCT	CTGAGAAATA	GGTGAGGCGA	AACAAAAAAT	9720
AATTTGCATG	ATAAAAAGTC	TGATTTTATG	ATCGCTATCC	TCGCTCTCTG	AGAAAGAAGC	9780
GAAACAAAGG	CGACTCCTGG	ACTCGAATCT	ATAAGATAAC	AAAGGCGACT	CCTGGGACTC	9840
GAATCTATAA	GATAACAAAG	GCAATTCCAA	GACTTGAATC	TATAAAAAAT	TTAGTTAAGA	9900
ATGATTAACG	TCCGATCCTA	ATTCGAATCG	AGGCATCTTA	CCACTCCATT	GATAATTATA	9960
TAAGTCAATA	AGTCATATAA	WAGTATTAAA	AACTAAATTG	ACTTGATCGG	TCTATCAAAA	10020
ATMAGATMAA	ATTGTGTTCA	TATGTAACAT	TTTTGTTGTC	ACAATTAGCT	TAATTACATC	10080
TTTCATGTGC	AATAACAAAG	AAATGATAGG	AATTTAGAGA	TTCCAATTTT	TTTGTTGCCA	10140
CAATTAACCT	AATTACATCT	TTCATTTGCA	ATAACAAAGA	AATGATAGGA	ATTTAGAGAT	10200
CCAGTGTCAA	TACACAACCT	AGGCCAACAT	CGAAAGCATA	ACTGTAAACT	CATGCATGAA	10260
GAAATCAGTC	GTA AAAATGA	ATAAATGCGA	CATAAAAACA	AATTGCATGT	ATCATTAATG	10320
TGACTTAACT	ACAAGTAAAA	ATAAATTTAA	CAAATGTAAC	TTAACTACAA	GTAAAAATAA	10380
ATTGCTTCTA	TCATTAACAA	ACAAACAGAA	TTAAAAAGAA	AAAAACATAC	TAAATCTTAC	10440
CGTCATTGCA	TAAAAAATAA	TACCAAATTC	ATAATGCAAG	GAAAACGAAA	CGCGTCCTGA	10500
TCGGGTATCA	ACGATGAAAT	GGACCAGTTG	GATCGACTGC	CTGCACAACG	TTAGGTATGC	10560
CAAAAAAAG	AACACGATCC	TTTGACCCCG	TTGATGATT	ATCAGTATGT	TCACAAAAAA	10620

AACTTAAGTT	CATCCCAGTG	TACAACAGCC	CCAACATCTG	CCCCAAGTAA	CAAAAAACAA	10680
CCAATTTATC	TTATTCTTAT	CTGCCACAAA	ATAATCGGTT	TCACACTATT	CTCTTGTTAT	10740
ACAAAATTGA	CAAGTAGGAA	GGAGAGGAGT	CATCCAAATA	AACGGTGCAC	GTTCTTTGAG	10800
AAAAGTCTTA	TTTTTCGTAA	GATCCAATTT	CAACAACTT	TTCTTCAAGT	CAAAATTCCT	10860
GATAGTGTAT	CTCCTCTCGA	CGACCTCTTG	CATTGAACGA	TCTCCGCTTA	TCATGAAAAG	10920
TTGCTTGGAT	AACAAGTATT	GCAAGGGGGG	GACAGTAGCT	ATTAAGTTAG	TCGGCCCAAG	10980
GAAATGGAGG	AGTGATAGTC	TCGAATATTA	TTCACCTCTT	TAGCATTACC	CGGTCTGGCT	11040
TTAAGGAGTT	ACGTCTTTTA	CGCTCGCCAA	TTTCTTTTTT	TAGAATGGTT	GGTGTCAAAA	11100
TCGCGAGTTG	TGGAAGGTTT	AAGTTACTCG	ATTTCGTGATT	TTCAAGTATG	AGTGGTGAGA	11160
GAGATTCGAT	ATTTTCACGA	GGTGTATTCT	AGGTCTAGTA	GAACGAAGGG	TGTCACTAAT	11220
GAAAGTTTCA	AGAGTTCATC	ATCATCTTCT	TCTAGTAGAT	TTTCGCTTTC	AAATGAGTAT	11280
GAAAATTCTT	CCTCTTTTCT	ATTGATTTTC	TTCATTGTTT	TCTTCATTGT	TGTGGTTGTT	11340
ATTGAAAAGA	AAGAAAATTT	ATAACAGAAA	AAGATGTCAA	AAAAAAGGTA	AAATGAAAGA	11400
GTATCATATA	CTTAAAGAGT	TGCGTAGAGA	TAAGTCAAAA	GAAACAGAAT	TATAGTAATT	11460
TCAGCTAAGT	TAGAATTC					11478

CLAIMS

1. A method of affecting enzymatic activity in a plant (or a cell, a tissue or an organ thereof) comprising expressing in the plant (or a cell, a tissue or an organ thereof) a nucleotide sequence wherein the nucleotide sequence codes, partially or completely, for an intron in a sense orientation; and wherein the nucleotide sequence does not contain a sequence that is sense to an exon sequence normally associated with the intron.

5
2. A method according to claim 1 wherein starch branching enzyme activity is affected and/or wherein the levels of amylopectin are affected and/or the composition of starch is changed.

10
3. A method of affecting enzymatic activity in a starch producing organism (or a cell, a tissue or an organ thereof) comprising expressing in the starch producing organism (or a cell, a tissue or an organ thereof) a nucleotide sequence wherein the nucleotide sequence codes, partially or completely, for an intron in a sense orientation; wherein the nucleotide sequence does not contain a sequence that is sense to an exon sequence normally associated with the intron; and wherein starch branching enzyme activity is affected and/or the levels of amylopectin are affected and/or the composition of starch is changed.

15

20
4. A method according to any one of claims 1 to 3 wherein the nucleotide sequence does not contain a sequence that is sense to an exon sequence.


25
5. A method according to any one of the preceding claims wherein the enzymatic activity is reduced or eliminated.
6. A method according to any one of the preceding claims wherein the nucleotide sequence codes for at least substantially all of at least one intron in a sense orientation.

30


7. A method according to any one of the preceding claims wherein the nucleotide sequence codes for all of at least one intron in a sense orientation.
8. A method according to any one of the preceding claims wherein the nucleotide sequence comprises the sequence shown as any one of SEQ.I.D. No. 1 to SEQ.I.D. No. 13 or a variant, derivative or homologue thereof, including combinations thereof.
9. A method according to any one of the preceding claims wherein the nucleotide sequence is expressed by a promoter having a sequence shown as SEQ.I.D. No. 14 or a variant, derivative or homologue thereof.
10. A sense sequence comprising the nucleotide sequence as defined in claim 8 or a variant, derivative or homologue thereof.
11. A promoter having a sequence shown as SEQ.I.D. No. 14, or a variant, derivative or homologue thereof.
12. A promoter according to claim 11 in combination with a gene of interest ("GOI").
13. A construct capable of comprising or expressing the invention according to any one of claims 10 to 12.
14. A vector comprising or expressing the invention according to any one of claims 10 to 13.
15. A combination of nucleotide sequences comprising a first nucleotide sequence coding for a recombinant enzyme; and a second nucleotide sequence which corresponds to an intron in a sense orientation; wherein the intron is an intron that is associated with a genomic gene encoding an enzyme corresponding to the recombinant enzyme; and wherein the second nucleotide sequence does not contain a sequence that is sense to an exon sequence normally associated with the intron.

16. A cell, tissue or organ comprising or expressing the invention according to any one of claims 10 to 15.
- 5 17. A transgenic starch producing organism comprising or expressing the invention according to any one of claims 10 to 16.
18. A transgenic starch producing organism according to claim 17 wherein the organism is a plant.
- 10 19. A starch obtained from the invention according to any one of the preceding claims.
20. pBEA11 (NCIMB 40754).
- 15 21. An intron nucleotide sequence that is obtainable from λ -SBE 3.2 (NCIMB 40751) or λ -SBE 3.4 (NCIMB 40752) or a variant, derivative or homologue thereof.
- 20 22. A method of expressing a recombinant protein or enzyme in a host organism comprising expressing a nucleotide sequence coding for the recombinant protein or enzyme; and expressing a further nucleotide sequence; wherein the further nucleotide sequence codes, partially or completely, for an intron in a sense orientation; wherein the intron is an intron normally associated with the genomic gene encoding a protein or an enzyme corresponding to the recombinant protein or enzyme; and wherein the further nucleotide sequence does not contain a sequence that is sense to an exon
- 25 sequence normally associated with the intron.

1/17



Reducing end



Reducing end




Fig 1

2/17




Fig 2




Fig 3

4/17




Fig 4

5/17




Fig 5

6/17




Fig 6

7/17




Fig 7

8/17

10	20	30	40	50	60	
123456789012345678901234567890123456789012345678901234567890						
ATCATGGCCAATTACTGGTTCAAATGCATTACTTCCTTTCAGATTCTTTCGAGTTCTCAT						60
GACCGGTCCTACTACAGACGATACTAACCCGTGGAAGTGTGCATCTGCTTCTTAGAACT						120
CTATGGCTATTTTCGTTAGCTTGGCGTCGGTTTGAACATAGTTTTTGTTCCTCAACTCTT						180
CATTTACAGTCAAAATGTTGTATGGTTTTTGTTCCTCAATGATGTTTACAGTGTGTG						240
TTGTCATCTGTACTTTTGCCTATTACTTGTTCCTCAATGATGTTTACAGTGTGTG						300
TTGCCATATTTTGTTCCTTATTATTATTATCATAACATATTATTACAAGGAAAAGACA						360
AGTACACAGATCTTAACGTTTATGTTCAATCAACTTTTGGAGGCATTGACAGGTACCACA						420
AATTTTGAGTTTATGATTAAGTTCAATCTTAGAATATGAATTTAACATCTATTATAGATG						480
CATAAAAATAGCTAATGATAGAACATTGACATTTGGCAGAGCTTAGGGTATGGTATATCC						540
AACGTTAATTTAGTAATTTTGTACGTACGTATATGAAATATTGAATTAATCACATGAA						600
CGGTGGATATTATATTATGAGTTGGCATCAGCAAAATCATTGGTGTAGTTGACTGTAGTT						660
GCAGATTTAATAATAAAATGGTAATTAACGGTCGATATTAAAATAACTCTCATTTCAGT						720
GGGATTAGAACTAGTTATTAAAAAATGTATACTTTAAGTGATTTGATGGCATATAATTT						780
AAAGTTTTTCATTTTCATGCTAAAATTGTTAATTATTGTAATGTAGACTGCGACTGGAATT						840
ATTATAGTGTAATTTATGCATTCAGTGTAATAATTAAAGTATTGAACCTGTCTGTTTTAG						900
AAAATACTTTTATCTTTAATATAGGATTTTGTTCATGCGAATTTAAATTAATCGATATTGA						960
ACACGGAATACCAAAATTAAAAAGGATACACATGGCCTTCATATGAACCGTGAACCTTTG						1020
ATAACGTGGAAGTTCAAAGAAGGTAAAGTTTAAAGTAATAAAGTGAACAAATTAATTTCTTTT						1080
ATTTGGCCCACTACTAAATTTGCTTTACTTTCTAACATGTCAAGTTGTGCCCTCTTAGTT						1140
GAATGATATTCATTTTTCATCCCATAAGTTCAATTTGATTGTGCATACCACCCATGATGTT						1200
CTGAAAAATGCTTGGCCATTACAAAGTTTATCTTAGTTTCTATGAACCTTTATAAGAAGC						1260
TTTAATTTGACATGTTATTATATTAGATGATATAATCCATGACCCAATAGACAAGTGTA						1320
TTAATATTGTAACCTTTGTAATTGAGTGTGTCTACATCTTATTCAATCATTTAAGGTCATT						1380
AAAATAAATTTATTTTGGACATTCTAAAACCTTAAGCAGAATAAATAGTTTATCAATTAT						1440
TAAAAACAAAAACGACTTATTTATAAATCAACAAACAATTTTAGATTGCTCCAACATAT						1500

Fig 8

9/17

10	20	30	40	50	60	
1234567890123456789012345678901234567890123456789012345678901234567890						
TTTTCCAAATTAAATGCAGAAAATGCATAATTTTATACTTGATCTTTATAGCTTATTTTT						1560
TTTAGCCTAACCAACGAATATTTGTAAACTCACAACCTTGATTAAAAGGATTACAAACAA						1620
GATATATATAAGTAGTGACAAATCTTGATTTTAAATATTTTAATTTGGAGGTCAAAATTT						1680
TACCATAATCATTTGTATTTATAATTAAATTTTAAATATCTTATTTATACATATCTAGTA						1740
AACTTTTAAATATACGTATATACAAAATATAAAATTATTGGCGTTCATATTAGGTCAATA						1800
AATCCTTAACTATATCTGCCTTACCACTAGGAGAAAGTAAAAAACTCTTTACCAAAAATA						1860
CATGTATTATGTATACAAAAGTCGATTAGATTACCTAAATAGAAATTGTATAACGAGTA						1920
AGTAAGTAGAAATATAAAAAAACTACAATACTAAAAAAATATGTTTTACTTCAATTTTCG						1980
AAACTAATGGGGTCTGAGTGAAATATTTCAGAAAGGGGAGGACTAACAAAAGGGTCATAAT						2040
GTTTTTTTATAAAAAGCCACTAAAATGAGGAAATCAAGAATCAGAACATACAAGAAGGCA						2100
GCAGCTGAAGCAAAGTACCATAATTTAATCAATGGAAATTAATTTCAAAGTTTATCAAA						2160
ACCCATTCGAGGATCTTTTCCATCTTTCTCACCTAAAGTTTCTTCAGGGGtaatttttac						2220
P I R G S F P S F S P K V S S G						
taatttcatgttaatttcaattattttttagcctttgcattttccaatatatctgg						2280
atcatctccttagttttttatttttatatttttataatatcaaatatggaagaaaaatgaca						2340
cttgtagagccatatgtaagtatcatgtgacaaatttgcaaggtggttgagtgtataaaa						2400
ttcaaaaattgagagatggaggggggggtgggggbaragacaatattagaaagagtgttc						2460
taggaggttatggaggacacggatgaggggtagaagggttaggttaggtatttgagtgttgt						2520
ctggcttatcctttcatactagtagtcgtggaattatttgggtagtttcttggtttgtta						2580
tttgatctttgttattctattttctgtttcttgacttcgattattgtattatatatctt						2640
gtcgtagttattgttcctcggttaagaatgctctagcatgcttccttagtgttttatcat						2700
gccttctttatattcgcggttgctttgaaatgcttttacttttagccgaggggtctattagaa						2760
acaatctctctatctcgtaaggtaggggttaaagtcctcaccacactccacttggtgggatt						2820
acattgtgtttgttggtgttaaataattatgtatacataataagtggattttttacaaca						2880
caaatacatggtcaagggcaaagttctgaacacataaaggggttcattatatgtccagga						2940
tatgataaaaattgtttctttgtgaaagttatataagatttggttatggcttttgctggaa						3000

Fig 8 continued.

10/17

10	20	30	40	50	60	
12345678901234567890123456789012345678901234567890						
acataataagttataatgctgagatagctactgaagtttgttttttctagccttttaaat						3060
gtaccaataatagattccgtatcgaaacgagtatgttttgattacctgggtcatgatgtttc						3120
tattttttacatttttttgggtgttgaaactgcaattgaaaatgttgatcctatgagacgg						3180
atagttgagaatgtgttctttgtatggaccttgagaagctcaaacgctactccaataatt						3240
tctatgaattcaaattcagtttatggctaccagtcagtcagaaattaggatatgctgca						3300
tatacttgttcaattatactgtaaaatttcttaagttctcaagatatccatgtaacctcg						3360
agaatttctttgacagGCTTCTAGAAATAAGATATGTTTTCTTCTCAACATAGTACTGG						3420
ACTGAAGTTTGGATCTCAGGAACGGTCTTGGGATATTTCTTCCACCCCAAAATCAAGAGT						3480
L K F G S Q E R S W D I S S T P K S R V						
TAGAAAAGATGAAAGGgtatgtttgataatttatatggttgcagtgatagatatataaata						3540
R K D E R						
gttggaaaacttctggactggtgctcatggcatatttgatctgtgcaccgtgtggagatg						3600
tcaaacatgtgttacttccgttccgccaatttataataccttaacttgggaaagacagctc						3660
tttactcctgtgggcatttgttatttgaattacaatctttatgagcatggtgttttcaca						3720
ttatcaacttctttcatgtggtatataacagtttttagctccgttaatacctttcttctt						3780
tttgatataaaactaactgtggtgcattgcttgcbkkkATGAAGCACAGTTCAGCTATTTT						3840
CGCTGTTTTGACCGATGACGACAATTCGACAATGGCACCCCTAGAGGAAGATGTCAAGAC						3900
A V L T D D D N S T M A P L E E D V K T						
TGAAAATATTGGCCTCCTAAATTTGGATCCAACCTTTGGAACCTTATCTAGATCACTTCAG						3960
E N I G L L N L D P T L E P Y L D H F R						
ACACAGAATGAAGAGATATGTGGATCAGAAAATGCTCATTGAAAAATATGAGGGACCCCT						4020
H R M K R Y V D Q K M L I E K Y E G P L						
TGAGGAATTTGCTCAAGgtaacagccaaaagttgtgctttaggcagtttgaccttatttt						4080
E E F A Q G						
ggaagatgaattgtttatacctactttgactttgctagagaattttgcataccggggagt						4140
aagtagtggctccatttaggtggcacctggccatttttttgatcttttaaaaagctgttt						4200
gattgggtcttcaaaaaagtagacaaggtttttggagaagtgcacacccccggagtgtc						4260
agtggcaaagcaaagattttcactaaggagattcaaaatataaaaaaagtatagacataa						4320
agaagctgaggggattcaacatgtactatacaagcatcaaatatagtcttaaagcaattt						4380
tgtagaaataaagaaagtcttccttctgttgcttcacaatttccttctattatcatgagt						4440
tactctttctgttcgaaatagcttccttaataattcatgatacttttgttgagatt						4500

Fig 8 continued

11/17

10	20	30	40	50	60	
123456789012345678901234567890123456789012345678901234567890						
tagcaggttttttcttgtgtaaactgctctctttttttgagGTTATTTAAATTTGGATT						4560
CAACAGGGAAGATGGTTGCATAGTCTATCGTGAATGGGCTCCTGCTCAGtaggtcct				Y L K F G F		4620
N R E D G C I V Y R E W A P A A Q						4680
cgtctactacaaaatagtagtttccatcatcataacagattttcctattaaagcatgatg						4740
ttgcagcatcattggctttcttacatgttctaattgctattaaggttatgcttctaatta						4800
actcatccacaatgcagGGAAGCAGAAGTTATTGGCGATTTCATGGATGGAACGGTTCT						4860
AACCACATGATGGAGAAGGACCAGTTTGGTGTGGAGTATTAGAATTCCTGATGTTGAC						4920
N H M M E K D Q F G V W S I R I P D V D						4980
AGTAAGCCAGTCATTCCACACAACCTCCAGAGTTAAGTTTCGTTTCAAACATGGTAATGGA						5040
S K P V I P H N S R V K F R F K H G N G						5100
GTGTGGGTAGATCGTATCCCTGCTTGGATAAAGTATGCCACTGCAGACGCCACAAAGTTT						5160
V W V D R I P A W I K Y A T A D A T K F						5220
GCAGCACCATATGATGGTGTCTACTGGGACCCACCCTTCAGAAAGgttttgttattca						5280
A A P Y D G V Y W D P P P S E R						5340
taccttgaagctgaattttgaacaccatcatcacaggcatttcgattcatgttcttacta						5400
gtcttgttatgtaagacattttgaaatgcaaaaggttaaataattgtgtctttactaatt						5460
tggacttgatcccatactctttcccttaacaaaatgagtcaattctataagtgcttgaga						5520
acttactacttcagcaattaaacagGTACCACTTCAAATACCCCTCGCCCTCCCAAACCCC						5580
GAGCCCCACGAATCTATGAAGCACATGTCGGCATGAGCAGCTCTGAGCCACGTGTAAATT						5640
A P R I Y E A H V G M S S S S E P R V N S						5700
CGTATCGTGAGTTTGCAGATGATGTTTTACCTCGGATTAAGGCAAATAACTATAACTG						5760
Y R E F A D D V L P R I K A N N Y N T V						5820
TCCAGTTGATGGCCATAATGGAACATTCTTACTATGGATCATTGGATATCATGTTACAA						5880
Q L M A I M E H S Y Y G S F G Y H V T N						5940
ACTTTTTTCTGTGAGCAGTAGATATGGAACCCGGAGGACCTAAAGTATCTGATAGATA						6000
F F A V S S R Y G N P E D L K Y L I D K						
AAGCACATAGCTTGGGTTTACAGGTTCTGGTGGATGTAGTTCACAGTCATGCAAGCAATA						
A H S L G L Q V L V D V V H S H A S N N						
ATGCACTGATGGCCTCAATGGCTTTGATATTGGCCAAGGTTCTCAAGAATCCTACTTTC						
V T D G L N G F D I G Q G S Q E S Y F H						
ATGCTGGAGAGCGAGGGTACCATAAGTTGTGGGATAGCAGGCTGTTCAACTATGCCAATT						
A G E R G Y H K L W D S R L F N Y A N W						
GGGAGGTTCTTCGTTTCCTTCTTTCCAACCTTGAGGTGGTGGCTAGAAGAGTATAACTTTG						
E V L R F L L S N L R W W L E E Y N F D						
ACGGATTTGATTTGATGGAATAACTTCTATGCTGTATGTTTCATCATGGAATCAATATGG						
G F R F D G I T S M L Y V H H G I N M G						
GATTTACAGGAACTATAATGAGTATTTACAGGAGGCTACAGATGTTGATGCTGTGGTCT						
F T G N Y N E Y F S E A T D V D A V V Y						
ATTTAATGTTGGCCAATAATCTGATTCACAAGATTTTCCCAGATGCAACTGTTATTGCCG						
L M L A N N L I H K I F P D A T V I A E						
AAGATGTTTCTGGTATGCCGGCCTTGGCCGGCTGTTTCTGAGGGAGGAATTGGTTTGG						
D V S G M P G L G R P V S E G G I G F V						

Fig 8 continued

12/17

10	20	30	40	50	60	
12345678901234567890123456789012345678901234567890						
TTTACCGCCTGGCAATGGCAATCCCAGATAAGTGGATAGATTATTTAAAGAATAAGAATG						6060
Y R L A M A I P D K W I D Y L K N K N D						
ATGAAGATTGGTCCATGAAGGAAGTAACATCGAGTTTGACAAATAGGAGATATACAGAGA						6120
E D W S M K E V T S S L T N R R Y T E K						
AGTGTATAGCATATGCGGAGACCCATGATCAGgtatttttaaatttatttctacaactaaa						6180
C I A Y A E T H D Q						
taatttctcagaacaattgtagatagaatccaaatatatcgtcctgaaagtataaaaagt						6240
acttatttttcgcatgggcttcagaatattggtagccgctgaatatcatgataagttat						6300
ttatccagtgacatttttatgttccactcctattatgtctgctggatacagTCTATTGTTG						6360
GTGACAAGACCATTGCAATTTCTCCTAATGGACAAAGAGATGTATTCTGGCATGTCTTGCT				S I V G		
D K T I A F L L M D K E M Y S G M S C L						6420
TGACAGATGCTTCTCTGTTGTTGATCGAGGAATTGCGCTTCACAAGgtttgtctgtttc						6480
T D A S P V V D R G I A L H K						
tattgcattttaaggttcatatagggttagccacggaaaatctcactctttgtgaggtaac						6540
caggggtctgatggattatttcaattttctcgttttatcatttgtttattcttttcatgcat						6600
tgtgtttctttttcaatatccctcttatttggaggtaattttttctcatctattcactttt						6660
agcttcttaaccacagATGATCCATTTTTCACAATGGCCTTGGGAGGAGAGGGGTACCTC						6720
M I H F F T M A L G G E G Y L						
AATTTTCATGGGTAACGAGgtatgtcttacatcttttagatattttgtgataattacaatta						6780
N F M G N E						
gtttggcttacttgaacaagattcattcctcaaaatgacctgaactggttgaacatcaaag						6840
gggttgaaacatagaggaaaacaacatgatgaatgtttccattgtctagggatttctatt						6900
atgttgctgagaacaaatgtcatcttaaaaaaacattgtttactttttttagtagataga						6960
agattactgtatagagtttgcaagtgtgtctgttttggagtaattgtgaaatgtttgatg						7020
aacttgtacagTTTGGCCATCCTGAGTGGATTGACTTCCCTAGAGAGGGCAATAATTGGA						7080
F G H P E W I D F P R E G N N W S						
GTTATGACAAATGTAGACGCCAGTGAACCTCGCGGATAGCGAACACTTGAGATACAAGg						7140
Y D K C R R Q W N L A D S E H L R Y K						
ttcaagtattttgaatcgagcttgttaaataatctagtaatttttagattgcttacttg						7200
gaagtctacttggttctggggatgatagctcatttcattctgttctacttattttccaac						7260
cgaatttctgatttttgtttcgagatccaagtattagattcatttacacttattaccgcc						7320
tcatttctaccactaaggccttgatgagcagcttaagttgattctttgaagctatagttt						7380
caggctaccaatccacagcctgctatatattgttggatacttaccttttctttacaatgaa						7440
gtgatactaattgaaatggtctaaatctgatattctatatttctccgtctttcctccccct						7500

Fig 8 continued

13/17

10	20	30	40	50	60	
12345678901234567890123456789012345678901234567890						
catgatgaaatgcagTTATGAATGCATTGATAGAGCTATGAATTCGCTCGATGAAAAG						7560
	F M N A F D R A M N S L D E K					
TTCTCATTCTCGCATCAGGAAAACAGATAGTAAGCAGCATGGATGATGATAATAAGgta						7620
F S F L A S G K Q I V S S M D D D N K						
aaatcatctaaagtgaaagtggtgggtttatgaagtgctttaattctatccaaggacaa						7680
gtagaaacctttttaccttccatttcttgatgatggatttcatattatttaatccaatag						7740
ctgggtcaaattcggtaatagctgtactgatttagttacttcactttgcagGTTGTTGTGTT						7800
	V V V F					
TGAACGTGGTGACCTGGTATTTGTATTCAACTTCCACCCAAAGAACACATACGAAGGgta						7860
E R G D L V F V F N F H P K N T Y E G						
tatatgttttacttatccatgaaattattgctctgcttggttttaattgtactgaacaagt						7920
tttatggagaagtaactgaaacaaatcattttcacattgtctaatttaactcttttttct						7980
gatcctcgcatgacgaaaacagGTATAAAGTTGGATGTGACTTGCCAGGGAAGTACAGAG						8040
	Y K V G C D L P G K Y R V					
TTGCACTGGACAGTGATGCTTGGGAATTTGGTGGCCATGGAAGgtaaggatttgcttga						8100
A L D S D A W E F G G H G R						
ataacttttgataataagataacagatgtagggtacagttctctcaccaaaaagaactgt						8160
aattgtctcatccatcttttagttgtataagatatccgactgtctgagttcggaagtgttt						8220
gagcctcctgccctccccctgcgttggttagctaattcaaaaaggagaaaactgtttatt						8280
gatgatctttgtcttcatgctgacatacaatctgttctcatgacagACTGGTCATGATGT						8340
	T G H D V					
TGACCATTTCACATCACCAGAAGGAATACCTGGAGTTCCAGAAACAAATTTCAATGGTCG						8400
D H F T S P E G I P G V P E T N F N G R						
TCCAAATTCCTTCAAAGTGCTGTCTCCTGCGCGAACATGTGTGgtacagttcttgccgtg						8460
P N S F K V L S P A R T C V						
tgacctccctttttattgtgggtttgttcatagttatttgaatgcgatagaagttaacta						8520
ttgattaccgccacaatcgccagtttaagtcctctgaactactaatttgaaaggtaggaat						8580
agccgtaataagggtctacttttggcatcttactgttacaaaacaaaaggatgccaaaaaa						8640
attcttctctatcctctttttccctaaaccagtgcatgtagcttgacactgcataaactt						8700
aggtaaatgatcaaaaatgaagttgatgggaacttaaaaccgccctgaagtaaagctagg						8760
aatagtcataataatgtccacctttggtgtctgcgctaacatcaacaacaacatacctcgt						8820
gtagtcccacaaagtggtttcagggggagggttagagtgtatgcaaaacttactcctatct						8880
cagaggtagagaggattttttcaatagacccttggtcaagaaaaaaagtcacaaaaagaa						8940
gtaacagaagtgaaagcaacatgtgtagctaaagcgacccaacttggttgggactgaagt						9000

Fig 8 continued

14/17

10	20	30	40	50	60	
123456789012345678901234567890123456789012345678901234567890						
agttggtggtggtgaaacagtgc	atgtagatgaacacatgtcagaaaatggacaacacag					9060
ttat	ttt	gtgcaagtcaaaaaatgtactactat	ttt	ctt	gtgcagctttatgtatagaa	9120
aag	tt	aaataactaatgaat	ttt	gtctagcagaaaaatagcttggagagaaat	tttttata	9180
ttga	actaagctaactatattcatct	ttt	ctttttt	gtcttcttctctcctt	gtttgtgaag	9240
GCTTATTACAGAGTTGATGAACGCATGTCAGAACTGAAGATTACCAGACAGACATTTGT						9300
A Y Y R V D E R M S E T E D Y Q T D I C						
AGTGAGCTACTACCAACAGCCAATATCGAGGAGAGTGACGAGAACTTAAAGATTCGTTA						9360
S E L L P T A N I E E S D E K L K D S L						
TCTACAAATATCAGTAACATTGACGAACGCATGTCAGAACTGAAGTTTACCAGACAGAC						9420
S T N I S N I D E R M S E T E V Y Q T D						
ATTTCTAGTGAGCTACTACCAACAGCCAATATTGAGGAGAGTGACGAGAACTTAAAGAT						9480
I S S E L L P T A N I E E S D E K L K D						
TCGTTATCTACAAATATCAGTAACATTGATCAGACTGTTGTAGTTTCTGTTGAGGAGAGA						9540
S L S T N I S N I D Q T V V V S V E E R						
GACAAGGAACCTTAAAGATTCACCGTCTGTAAGCATCATTAGTGATGTTGTTCCAGCTGAA						9600
D K E L K D S P S V S I I S D V V P A E						
TGGGATGATTGAGATGCAACGCTCTGGGGTGAGGACTAGTCAGATGATTGATCGACCCTT						9660
W D D S D A N V W G E D						
CTACCGATTGGTGATCGCTATCCTTGCTCTCTGAGAAATAGGTGAGGCGAAACAAAAAAT						9720
AATTTGCATGATAAAAAAGTCTGATTTTTATGATCGCTATCCTCGCTCTCTGAGAAAGAAGC						9780
GAAACAAAGGCGACTCCTGGACTCGAATCTATAAGATAACAAAGGCGACTCCTGGGACTC						9840
GAATCTATAAGATAACAAAGGCAATTCCAAGACTTGAATCTATAAAAAATTTAGTTAAGA						9900
ATGATTAACGTCCGATCCTAATTCGAATCGAGGCATCTTACCACTCCATTGATAATTATA						9960
TAAGTCAATAAGTCATATAAWAGTATTAAAAACTAAATTGACTTGATCGGTCTATCAAAA						10020
ATMAGATMAAATTGTGTTTCATATGTAACATTTTGTGTTGTCACAATTAGCTTAATTACATC						10080
TTTCATGTGCAATAACAAAGAAATGATAGGAATTTAGAGATTCCAATTTTTTTGTTGCCA						10140
CAATTAACCTTAATTACATCTTTTCATTTGCAATAACAAAGAAATGATAGGAATTTAGAGAT						10200
CCAGTGTCATACACAACCTAGGCCAACATCGAAAGCATAACTGTAAACTCATGCATGAA						10260
GAAATCAGTCGTAAAAATGAATAAATGCGACATAAAAAACAAATTGCATGTATCATTAAATG						10320
TGACTTAACTACAAGTAAAAATAAATTTAACAAATGTAACCTAACTACAAGTAAAAATAA						10380
ATTGCTTCTATCATTAACAAACAAACAGAATTAAGAAAGAAAAAACATACTAAATCTTAC						10440
CGTCATTTCGATAAAAAAAATACCAAATTCATAATGCAAGGAAAACGAAACGCGTCTGA						10500

Fig 8 continued

15/17

10	20	30	40	50	60	
12345678901234567890123456789012345678901234567890						
TCGGGTATCAACGATGAAATGGACCAGTTGGATCGACTGCCTGCACAACGTTAGGTATGC						10560
CAAAAAAAGAACACGATCCTTTGCACCCGTTTCGATGATTATCAGTATGTTCAAAAAA						10620
AACTTAAGTTCATCCCAGTGTAACAACAGCCCCAACATCTGCCCCAAGTAACAAAAACAA						10680
CCAATTTATCTTATTCTTATCTGCCACAAAATAATCGGTTTCACACTATTCTCTTGTAT						10740
ACAAAATTGACAAGTAGGAAGGAGAGGAGTCATCCAAATAAACGGTGACGTTCTTTGAG						10800
AAAAGTCTTATTTTTCGTAAGATCCAATTTCAACAACTTTTCTTCAAGTCAAAATTCCT						10860
GATAGTGTATCTCCTCTCGACGACCTCTTCGATTGAACGATCTCCGCTTATCATGAAAAG						10920
TTGCTTGGATAACAAGTATTGCAAGGGGGGACAGTAGCTATTAAGTTAGTCGGCCCAAG						10980
GAAATGGAGGAGTGATAGTCTCGAATATTATTACCTCTTTAGCATTACCCGGTCTGGCT						11040
TTAAGGAGTTACGTCTTTTACGCTCGCCAATTTCTTTTTTTAGAAATGGTTGGTGTCAAAA						11100
TCGCGAGTTGTGGAAGGTTCAAGTTACTCGATTTCGTGATTTTCAAGTATGAGTGGTGAGA						11160
GAGATTCGATATTTTCACGAGGTGTATTCGAGGTCTAGTAGAACGAAGGGTGTCACTAAT						11220
GAAAGTTTCAAGAGTTCATCATCATCTTCTTCTAGTAGATTTTCGCTTTCAAATGAGTAT						11280
GAAAATCTTCCTCTTTTCTATTGATTTTCTTCATTGTTTCTTCATTGTTGTGGTTGTT						11340
ATTGAAAAGAAAGAAATTTATAACAGAAAAAGATGTCAAAAAAAGGTAAATGAAAGA						11400
GTATCATATACTTAAAGAGTTGCGTAGAGATAAGTCAAAAGAAACAGAATTATAGTAATT						11460
TCAGCTAAGTTAGAATTC						11478

Fig 8 continued

16/17




Fig 9

17/17





Fig 10

PLT/EP 96 / 03053

INDICATIONS RELATING TO A DEPOSITED MICROORGANISM

(PCT Rule 13bis)

A. The indications made below relate to the microorganism referred to in the description on page <u>19</u> , lines <u>18 to 27</u>	
B. IDENTIFICATION OF DEPOSIT Further deposits are identified on an additional sheet <input type="checkbox"/>	
Name of depositary institution The National Collections of Industrial and Marine Bacteria Limited (NCIMB)	
Address of depositary institution (including postal code and country) 23 St. Machar Drive Aberdeen Scotland AB2 1RY United Kingdom	
Date of deposit 13 July 1995	Accession Number NCIMB 40754, NCIMB 40751, NCIMB 40752
C. ADDITIONAL INDICATIONS (leave blank if not applicable) This information is continued on an additional sheet <input type="checkbox"/>	
In respect of those designations in which a European patent is sought, and any other designated state having equivalent legislation, a sample of the deposited microorganism will be made available until the publication of the mention of the grant of the European patent or until the date on which the application has been refused or withdrawn or is deemed to be withdrawn, only by the issue of such a sample to an expert nominated by the person requesting the sample. (Rule 28(4) EPC).	
D. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE (if the indications are not for all designated States)	
E. SEPARATE FURNISHING OF INDICATIONS (leave blank if not applicable)	
The indications listed below will be submitted to the International Bureau later (specify the general nature of the indications e.g., "Accession Number of Deposit")	
<div>For receiving Office use only</div> <div><input checked="" type="checkbox"/> This sheet was received with the international application</div> <div></div> <div>Authorized officer P.L.R. Pether</div>	<div>For International Bureau use only</div> <div><input type="checkbox"/> This sheet was received by the International Bureau on:</div> <div>Authorized officer</div>

BUDAPEST TREATY ON THE INTERNATIONAL
RECOGNITION OF THE DEPOSIT OF MICROORGANISMS
FOR THE PURPOSES OF PATENT PROCEDURE

PCT/EP 96 / 03053

Danisco Biotechnology
1. Artvejsgade 1
P O Box 17
DK-1001 Copenhagen K
Denmark

INTERNATIONAL FORM

RECEIPT IN THE CASE OF AN ORIGINAL DEPOSIT
issued pursuant to Rule 7.1 by the
INTERNATIONAL DEPOSITARY AUTHORITY
identified at the bottom of this page

I. IDENTIFICATION OF THE MICROORGANISM	
Identification reference given by the DEPOSITOR:	Accession number given by the INTERNATIONAL DEPOSITARY AUTHORITY:
Escherichia coli DH5 α -pBEA11	NCIMB 40754
II. SCIENTIFIC DESCRIPTION AND/OR PROPOSED TAXONOMIC DESIGNATION	
The microorganism identified under I above was accompanied by:	
<input type="checkbox"/> a scientific description	
<input checked="" type="checkbox"/> a proposed taxonomic designation	
(Mark with a cross where applicable)	
III. RECEIPT AND ACCEPTANCE	
This International Depositary Authority accepts the microorganism identified under I above, which was received by it on 13 July 1995 (date of the original deposit) ¹	
IV. RECEIPT OF REQUEST FOR CONVERSION	
The microorganism identified under I above was received by this International Depositary Authority on (date of the original deposit) and a request to convert the original deposit to a deposit under the Budapest Treaty was received by it on (date of receipt of request for conversion)	
V. INTERNATIONAL DEPOSITARY AUTHORITY	
Name: Address:	Signature(s) of person(s) having the power to represent the International Depositary Authority or of authorized official(s): Date: 21 July 1995
NCIMB Ltd 23 St Machar Drive Aberdeen Scotland UK AB2 1RY	<i>Terence Dando</i>

¹ Where Rule 6.4(d) applies, such date is the date on which the status of international depositary authority was acquired.

BUDAPEST TREATY ON THE INTERNATIONAL
RECOGNITION OF THE DEPOSIT OF MICROORGANISMS
FOR THE PURPOSES OF PATENT PROCEDURE

Danisco Biotechnology
Langebrogade 1
P O Box 17
DK-1001 Copenhagen K
Denmark

INTERNATIONAL FORM

PCT/EP96/03053

VIABILITY STATEMENT
issued pursuant to Rule 10.2 by the
INTERNATIONAL DEPOSITARY AUTHORITY
identified on the following page

NAME AND ADDRESS OF THE PARTY
TO WHOM THE VIABILITY STATEMENT
IS ISSUED

I. DEPOSITOR	II. IDENTIFICATION OF THE MICROORGANISM
Name: As above Address:	Accession number given by the INTERNATIONAL DEPOSITARY AUTHORITY: NCIMB 40754 Date of the deposit or of the transfer: 13 July 1995
III. VIABILITY STATEMENT	
<p>The viability of the microorganism identified under II above was tested on 13 July 1995</p> <p>³ <input checked="" type="checkbox"/> viable</p> <p>³ <input type="checkbox"/> no longer viable</p>	

- ¹ Indicate the date of the original deposit or, where a new deposit or a transfer has been made, the most recent relevant date (date of the new deposit or date of the transfer).
- ² In the cases referred to in Rule 10.2(a)(ii) and (iii), refer to the most recent viability test.
- ³ Mark with a cross the applicable box.

I. CONDITIONS UNDER WHICH THE VIABILITY TEST HAS BEEN PERFORMED⁴

V. INTERNATIONAL DEPOSITARY AUTHORITY

Name:

NCIMB Ltd

Address:

23 St Machar Drive
Aberdeen Scotland
UK AB9 1RVSignature(s) of person(s) having the power
to represent the International Depositary
Authority or of authorized official(s):

Date:

Terence Dando
21 July 1995⁴ Fill in if the information has been requested and if the results of the test were negative.

BUDAPEST TREATY ON THE INTERNATIONAL
RECOGNITION OF THE DEPOSIT OF MICROORGANISMS
FOR THE PURPOSES OF PATENT PROCEDURE

DR A Buchter-Larsen
Danisco Biotechnology
Ingebrogade 1
P O Box 17
DK-1001 Copenhagen
Denmark

INTERNATIONAL FORM

PCT/EP 96 / 03053

RECEIPT IN THE CASE OF AN ORIGINAL DEPOSIT
issued pursuant to Rule 7.1 by the
INTERNATIONAL DEPOSITARY AUTHORITY
identified at the bottom of this page

I. IDENTIFICATION OF THE MICROORGANISM	
Identification reference given by the DEPOSITOR:	Accession number given by the INTERNATIONAL DEPOSITARY AUTHORITY:
Bacteriophage λ EMBL3 SP6/T7 λ SBE3.2	NCIMB 40751
II. SCIENTIFIC DESCRIPTION AND/OR PROPOSED TAXONOMIC DESIGNATION	
The microorganism identified under I above was accompanied by:	
<input type="checkbox"/> a scientific description	
<input checked="" type="checkbox"/> a proposed taxonomic designation	
(Mark with a cross where applicable)	
III. RECEIPT AND ACCEPTANCE	
This International Depositary Authority accepts the microorganism identified under I above, which was received by it on 13 July 1995 (date of the original deposit) ¹	
IV. RECEIPT OF REQUEST FOR CONVERSION	
The microorganism identified under I above was received by this International Depositary Authority on (date of the original deposit) and a request to convert the original deposit to a deposit under the Budapest Treaty was received by it on (date of receipt of request for conversion)	
V. INTERNATIONAL DEPOSITARY AUTHORITY	
Name: NCIMB Ltd 23 St Machar Drive Aberdeen Scotland Address: UK AB2 1FV	Signature(s) of person(s) having the power to represent the International Depositary Authority or of authorized official(s): Date: 20 July 1995

¹ Where Rule 6.4(d) applies, such date is the date on which the status of international depositary authority was acquired.

BUDAPEST TREATY ON THE INTERNATIONAL
RECOGNITION OF THE DEPOSIT OF MICROORGANISMS
FOR THE PURPOSES OF PATENT PROCEDURE

Dr A Buchter-Larsen
Danisco Biotechnology
Langebrogade 1
P O Box 17
DK-1001 Copenhagen K
Denmark

INTERNATIONAL FORM

VIABILITY STATEMENT
issued pursuant to Rule 10.2 by the
INTERNATIONAL DEPOSITARY AUTHORITY
identified on the following page

NAME AND ADDRESS OF THE PARTY
TO WHOM THE VIABILITY STATEMENT
IS ISSUED

I. DEPOSITOR	II. IDENTIFICATION OF THE MICROORGANISM
Name: As above Address:	Accession number given by the INTERNATIONAL DEPOSITARY AUTHORITY: NCIMB 40751 Date of the deposit or of the transfer: 13 July 1995
III. VIABILITY STATEMENT	
The viability of the microorganism identified under II above was tested on 19 July 1995 ² . On that date, the said microorganism was <div style="display: flex; justify-content: space-between; margin-top: 10px;"> <div data-bbox="162 1302 341 1365"> ³ <input checked="" type="checkbox"/> viable </div> <div data-bbox="162 1386 479 1449"> ³ <input type="checkbox"/> no longer viable </div> </div>	

¹ Indicate the date of the original deposit or, where a new deposit or a transfer has been made, the most recent relevant date (date of the new deposit or date of the transfer).

² In the cases referred to in Rule 10.2(a)(ii) and (iii), refer to the most recent viability test.

³ Mark with a cross the applicable box.

IV. CONDITIONS UNDER WHICH THE VIABILITY TEST HAS BEEN PERFORMED⁴

V. INTERNATIONAL DEPOSITARY AUTHORITY


Name:

Address:

NCIMB Ltd
23 St Machar Drive
Aberdeen Scotland
UK AB2 1RY

Signature(s) of person(s) having the power
to represent the International Depositary
Authority or of authorized official(s):

Date:


20 July 1995

⁴ Fill in if the information has been requested and if the results of the test were negative.

BUDAPEST TREATY ON THE INTERNATIONAL
RECOGNITION OF THE DEPOSIT OF MICROORGANISMS
FOR THE PURPOSES OF PATENT PROCEDURE

Dr A Buchter-Larsen
Danisco Biotechnology
Langebrogade 1
Box 17
DK-1001 Copenhagen
Denmark

INTERNATIONAL FORM

PCT/EP 96 / 03053

RECEIPT IN THE CASE OF AN ORIGINAL DEPOSIT
issued pursuant to Rule 7.1 by the
INTERNATIONAL DEPOSITARY AUTHORITY
identified at the bottom of this page

NAME AND ADDRESS
OF DEPOSITOR

I. IDENTIFICATION OF THE MICROORGANISM	
Identification reference given by the DEPOSITOR: Bacteriophage λ EMBL3 SP6/T7 λ SBE3.4	Accession number given by the INTERNATIONAL DEPOSITARY AUTHORITY: NCIMB 40752
II. SCIENTIFIC DESCRIPTION AND/OR PROPOSED TAXONOMIC DESIGNATION	
The microorganism identified under I above was accompanied by: <input type="checkbox"/> a scientific description <input checked="" type="checkbox"/> a proposed taxonomic designation (Mark with a cross where applicable)	
III. RECEIPT AND ACCEPTANCE	
This International Depositary Authority accepts the microorganism identified under I above, which was received by it on 13 July 1995 (date of the original deposit) ¹	
IV. RECEIPT OF REQUEST FOR CONVERSION	
The microorganism identified under I above was received by this International Depositary Authority on (date of the original deposit) and a request to convert the original deposit to a deposit under the Budapest Treaty was received by it on (date of receipt of request for conversion)	
V. INTERNATIONAL DEPOSITARY AUTHORITY	
Name: Address: NCIMB Ltd 23 St Machar Drive Aberdeen Scotland UK AR2 4YV	Signature(s) of person(s) having the power to represent the International Depositary Authority or of authorized official(s): Date: 20 July 1995

¹ Where Rule 6.4(d) applies, such date is the date on which the status of international depositary authority was acquired.

BUDAPEST TREATY ON THE INTERNATIONAL
RECOGNITION OF THE DEPOSIT OF MICROORGANISMS
FOR THE PURPOSES OF PATENT PROCEDURE

Dr A Buchter-Larsen
Danisco Biotechnology
Langebrogade 1
P O Box 17
DK-1001 Copenhagen K
Denmark

INTERNATIONAL FORM

VIABILITY STATEMENT
issued pursuant to Rule 10.2 by the
INTERNATIONAL DEPOSITARY AUTHORITY
identified on the following page

NAME AND ADDRESS OF THE PARTY
TO WHOM THE VIABILITY STATEMENT
IS ISSUED

I. DEPOSITOR	II. IDENTIFICATION OF THE MICROORGANISM
Name: As above Address:	Accession number given by the INTERNATIONAL DEPOSITARY AUTHORITY: NCIMB 40752 Date of the deposit or of the transfer: 13 July 1995
III. VIABILITY STATEMENT	
The viability of the microorganism identified under II above was tested on 19 July 1995 ¹ ² On that date, the said microorganism was <input checked="checked" type="checkbox"/> ³ viable <input type="checkbox"/> ³ no longer viable	

- ¹ Indicate the date of the original deposit or, where a new deposit or a transfer has been made, the most recent relevant date (date of the new deposit or date of the transfer).
- ² In the cases referred to in Rule 10.2(a)(ii) and (iii), refer to the most recent viability test.
- ³ Mark with a cross the applicable box.

IV. CONDITIONS UNDER WHICH THE VIABILITY TEST HAS BEEN PERFORMED⁴

V. INTERNATIONAL DEPOSITARY AUTHORITY


Name:

Address:

NCIMB Ltd
23 St Machar Drive
Aberdeen Scotland
AB9 1RY

Signature(s) of person(s) having the power
to represent the International Depositary
Authority or of authorized official(s):

Date:


20 July 1995

⁴ Fill in if the information has been requested and if the results of the test were negative.